

The Contribution of Leaching to the Rapid Release of Nutrients and Carbon in the Early

2 Decay of Wetland Vegetation

4 Stephen E. Davis, III^{1a}, Daniel L. Childers², and Gregory B. Noe³

6 ¹*Department of Wildlife & Fisheries Sciences, MS 2258, Texas A&M University, College Station, TX 77843; 979-458-3475 (voice), 979-845-4096 (fax), sedavis@tamu.edu*

8

²*Department of Biological Sciences & SE Environmental Research Center, Florida International University, Miami, FL 33199; 305-348-3101 (voice), 305-348-4096 (fax), childers@fiu.edu*

12 ³*U.S. Geological Survey, 430 National Center, Reston, VA 20192; 703-648-5826 (voice), 703-648-5484 (fax), gnoe@usgs.gov*

14

16 *Keywords: leaf decomposition, organic carbon, nitrogen, phosphorus, North Inlet, Everglades*

18 Note: This paper has not been submitted elsewhere in identical or similar form, nor will it be during the first three months after its submission to *Hydrobiologia*.

20

22 ^a Corresponding author

Abstract

2 Our goal was to quantify the coupled process of litter turnover and leaching as a source of
nutrients and fixed carbon in oligotrophic, nutrient-limited wetlands. We conducted poisoned
4 and non-poisoned incubations of leaf material from four different perennial wetland plants
(*Eleocharis* spp., *Cladium jamaicense*, *Rhizophora mangle*, and *Spartina alterniflora*) collected
6 from different oligotrophic freshwater and estuarine wetland settings. Total phosphorus (TP)
release from the P-limited Everglades plant species (*Eleocharis* spp., *C. jamaicense*, and *R.*
8 *mangle*) was much lower than TP release by the salt marsh plant *S. alterniflora* from N-limited
North Inlet (SC). For most species and sampling times, total organic carbon (TOC) and TP
10 leaching losses were much greater in poisoned than non-poisoned treatments, likely as a result of
epiphytic microbial activity. Therefore, a substantial portion of the C and P leached from these
12 wetland plant species was bio-available to microbial communities. Even the microbes associated
with *S. alterniflora* from N-limited North Inlet showed indications of P-limitation early in the
14 leaching process, as P was removed from the water column. Leaves of *R. mangle* released much
more TOC per gram of litter than the other species, likely contributing to the greater waterborne
16 [DOC] observed by others in the mangrove ecotone of Everglades National Park. Between the
two freshwater Everglades plants, *C. jamaicense* leached nearly twice as much P than *Eleocharis*
18 spp. In scaling this to the landscape level, our observed leaching losses combined with higher
litter production of *C. jamaicense* compared to *Eleocharis* spp. resulted in a substantially greater
20 P leaching from plant litter to the water column and epiphytic microbes. In conclusion, leaching
of fresh plant litter can be an important autochthonous source of nutrients in freshwater and
22 estuarine wetland ecosystems.

24

Introduction

2 Litter produced by deciduous and evergreen trees is the primary mechanism by which
nutrients are returned to the soil (Swift et al., 1979). Herbaceous plants also shed leaves and
4 other aboveground parts, contributing to the recycling of nutrients and organic matter (Teal,
1962; Moran & Hodson, 1989). This pool of litter represents a relatively large, labile reservoir
6 of organic matter to soil decomposer communities (Gosselink & Kirby, 1985; Benner et al.,
1986). Although resorption prior to leaf abscission can be an effective means of conserving vital
8 elements in many plant species, there is still a substantial outflow of organic and inorganic
nutrients from trees and macrophytes via leaf senescence and decomposition (Tukey, 1970; Gosz
10 et al., 1973; Aerts, 1996; Killingbeck, 1996; Maie et al., 2006).

The initial leaching phase of wood and leaf litter decomposition typically lasts from a few
12 days to a few weeks, yet it is responsible for substantial loss of mass and release of materials
such as carbon, nitrogen, and phosphorus to the environment (Ibrahima et al., 1995; Taylor &
14 Bärlocher, 1996; Davis et al., 2003; Romero et al., 2005). Given that standing water or saturated
soil conditions expedite this abiotic process (Tukey, 1970), the coupled process of litter
16 production and leaching may be an especially important autochthonous source of nutrients and
fixed carbon in oligotrophic, nutrient-limited wetland systems.

18 In order to address this, we conducted leaching studies on leaf material from four
different perennial wetland plants (*Eleocharis* spp., *Cladium jamaicense*, *Rhizophora mangle*,
20 and *Spartina alterniflora*) collected from different oligotrophic freshwater and estuarine wetland
settings. Our objectives were: 1) to quantify the time-dependent release of C, N, and P from
22 each of these species when immersed in water, 2) to evaluate the role of biological processes in
governing these physically-driven losses of C, N, and P from each species, and 3) using

published values on litter production in each of these wetland systems, we wanted to develop C, N, and P budgets for each species associated with early (i.e. < 3 weeks) decay. A common goal underlying these objectives was to identify links between oligotrophic status in wetland ecosystems and leaching as a mechanism for the internal recycling of both limiting and non-limiting elements.

Based on previous work by Davis et al. (2003), we hypothesized that leaching alone would be responsible for greater loss of materials than biotic processes (i.e. microbial degradation) during the first three weeks of decomposition. However, we expected the biological contributions to litter decay to increase over this period of time, as microbial colonization increased and leachable materials were exhausted. Lastly, given that these plants were all collected from oligotrophic, nutrient-limited systems, we anticipated that high resorption efficiency would result in would result in relatively low leachable fractions of limiting elements (Feller et al., 1999; Richardson et al., 1999). In other words, we expected that the quantity of phosphorus (normalized to dry mass of plant tissue) leached from phosphorus-limited plant tissue would be considerably less than phosphorus leached from a nitrogen-limited plant, and vice versa.

Materials and Methods

Between August 2000 to July 2002, we conducted leaching experiments on leaf material from four different wetland species. In 2000, we leached senesced, yellow leaves collected from an estuarine dwarf red mangrove (*Rhizophora mangle*) wetland along Taylor River, Everglades National Park, Florida USA (Table 1). In 2001, we leached senesced tissue from two freshwater macrophytes (*Cladium jamaicense* leaves and *Eleocharis* spp. culms) collected in southeast

Everglades National Park, just south of the C-111 canal (Table 1). Both Everglades wetlands
2 and associated plant communities are highly oligotrophic and limited by phosphorus availability
(Davis, 1989; Koch & Snedaker, 1997; Noe et al., 2001). Finally, in 2002, we leached senesced
4 leaf blades of *Spartina alterniflora* that were collected from North Inlet, South Carolina USA
(Table 1). Studies of this pristine salt marsh ecosystem indicate that macrophyte productivity is
6 strongly limited by the availability of nitrogen, although soil microbial processes are phosphorus
limited (Sundareshwar et al., 2003).

8 For all four species, only leaf material above the mean high water mark was collected,
which we assumed had not been significantly leached already. Leaves were air-dried from the
10 time of collection to the initiation of the experiment—approximately 48 hours later. Leaves
were weighed and incubated in 250 ml, clear, square, glass bottles containing 240 ml of water
12 from each site. For all four species, we used approximately 1–4 g air-dried leaf material in each
incubation bottle. Incubations lasted no longer than 21 days, as this is the timeframe needed to
14 fully capture the shift from abiotic to biotic contributions to decomposition (Davis et al., 2003).
We assumed that ambient surface water from each wetland would provide the most realistic,
16 wetland-specific environment for the early decay of each species. However, prior to each set of
incubations, we filtered (GF/F) the water to reduce variability in large particles ($> 0.7 \mu\text{m}$)
18 between different water sources. The source water for each set of incubations was fresh (i.e., 0
‰), except for the *S. alterniflora* incubation, which was mesohaline (15 ‰; Table 2).

20 To help distinguish the contribution of leaching from microbial processes in the early
phase (< 3 weeks) of leaf decomposition, we added a poison (2 ml of a 1% solution of NaN_3) to
22 half the bottles. The other half of the bottles received 2 ml of de-ionized water. All the bottles
from each study were incubated next to one another (spaced 2–5 cm apart from one another) in

an outdoor setting in shallow (< 0.5 m) water under ambient temperature and sunlight conditions.

2 All treatment combinations were conducted in triplicate for each set of incubations. In order to
understand how leaching losses changed over time, we sacrificed three “poisoned” and three
4 “non-poisoned” bottles after 1, 2, 5, 10 and 21 days of incubation. Studies have shown that
much of the leachable fraction is exhausted after 24 hours (Webster & Benfield, 1986).
6 However, for some components, leaching can be detected for several days (Ibrahima et al.,
1995).

8 During each sampling, leaves were removed from the bottles and water samples were
collected. Water samples were stored in 125 ml, HDPE bottles at 4°C until analyzed for C, N,
10 and P content. All water nutrient analyses were conducted at the Southeast Environmental
Research Center’s laboratory at Florida International University. Samples were analyzed for
12 total phosphorus (TP) according to a modification of the dry ashing, acid-hydrolysis technique
(Solorzano & Sharp, 1980), for total nitrogen (TN) using an Antec 7000N total nitrogen
14 analyzer, and for total organic carbon (TOC) using a hot platinum catalyst, direct injection
analyzer (Shimadzu model TOC-5000).

16 To ensure that changes in water nutrients were solely due to the leaves, control bottles
containing only water or water + poison were incubated for the entire 21-day length of each
18 experiment. Nutrient concentrations from the control bottles were compared with initial
concentrations to determine changes in C, N, and P fractions associated with water column or
20 photochemical processes. Paired t-tests were used to determine significant differences between
initial and final concentrations ($P < 0.05$). Since NaN_3 was selected as the poison in this
22 experiment, TN concentrations were more than an order of magnitude higher in the bottles

containing NaN_3 . Therefore, we are unable to report on the fluxes of TN in bottles containing
2 the poison.

Because we used air-dried leaf material, an accurate means of estimating initial oven-
4 dried mass was needed in order to normalize calculated releases of C, N, and P. To accomplish
this, we converted oven-dried mass to air-dried mass for each species (Table 1). These
6 conversions were generated from 25–30 individual leaves of each species that were weighed
after being air-dried during the same period of time as the experimental leaves (≤ 48 hours), then
8 oven-dried to a constant mass at 70°C . The conversion for each species involved multiplying by
the initial, air-dried mass in order to estimate initial dry mass for each experimental leaf.

10 The changes in nutrients in bottles containing leaf material and poison were assumed to
be the result of leaching. For those without poison, we assumed that both leaching and
12 biological processes were at work in governing C, N, and P dynamics. We calculated total
releases from each incubation bottle as the change in the molar quantities of TOC, TN, and TP
14 from initial source water, normalized to the predicted initial dry mass of leaf material in each
bottle per time of incubation (in moles \times gdw leaf material⁻¹ time⁻¹). We do not present data on
16 rates of C, N, and P release, but these values can be easily estimated by dividing the total release
by the number of days incubated (i.e., 1 or 21).

18 We used analysis of variance (ANOVA) to determine the effect of time on early decay of
leaf material, comparing the yield of C, N, and P leached from all species after 1 day of
20 incubation to total fluxes after 21 days of incubation. We also used ANOVA to determine the
effect that the poison had on releases of C, N, and P in each species to discern the time-
22 dependence of biological contributions to these releases. Lastly, we used ANOVA to determine
species effects on releases of each of these constituents over a three-week period of time. For

each of these analyses, Tukey-Kramer post-hoc tests were used to determine differences between
2 treatment means of significant ANOVAs ($P < 0.05$). By doing these analyses and focusing on
initial, 1-day releases and 21-day yield, we hoped to generate a better understanding the role of
4 microbes in governing the fate of leached C, N, and P for each species.

6 **Results**

All species released a significant amount of TOC, TN, and TP after just 1 day (Figures 1–
8 3), compared to control bottles (i.e., those without leaves) that showed no significant change in
these constituents over each of the 3-week experiments (see Table 2 for initial concentrations of
10 TOC, TN, and TP in each water source). We also collected C, N, and P data from each set of
incubations after 2, 5, and 10 days of incubation, but these data only followed the trends we
12 observed after 1 and 21 days. Therefore, we chose to omit these days from our discussion and
focus on the time end-members of these experiments.

14 For all species and poison treatment combinations, 21-day total releases of TOC and TP
were always significantly greater than 1-day total releases, indicating that a substantial pool of
16 leachable materials still existed within the leaves after one day (ANOVA; $p < 0.0001$).

However, the rate of release was highest in the first day for all species (ANOVA; $p < 0.0001$).

18 When comparing within species, *R. mangle* leaves released an order of magnitude more
TOC than the other three species regardless of the addition of poison (ANOVA; $p < 0.0001$).
20 From Day 1 to Day 21, the amount of TOC released by *R. mangle* jumped by more than ten-fold
(Figure 1). The other species did not exhibit this same magnitude of trend for TOC release, as
22 the increases from Day 1 to Day 21 shown by *C. jamaicense*, *Eleocharis spp.*, and *S. alterniflora*
were each less than three-fold (Figure 1). For these same three species, the difference between

poisoned and non-poisoned incubations was significant (poisoned > non-poisoned) during both days of sampling (Table 3), indicating a sustained biological effect on the early release of TOC.

Rhizophora mangle leaves revealed a similar biological effect after 1 day of immersion.

However, we could not statistically differentiate poisoned releases of TOC from non-poisoned releases by Day 21 in the mangrove leaf incubations (Table 3, Figure 1).

The amount of TP leached from these four species was considerably less than the amount of TOC released, ranging from sub- μ mole levels of TP (e.g., all Everglades species after 1 day of incubation) to as much as 13.5 μ moles TP gdw^{-1} from *S. alterniflora* blades after 21 days.

Overall, *Spartina alterniflora* blades released significantly more (by about an order of magnitude) TP than leaf material from the three Everglades species (ANOVA; $p < 0.0001$), which were limited by P availability in their respective natural settings. Total phosphorus leached from poisoned and non-poisoned incubations containing *R. mangle* leaves were not different after 1 day, but poisoned releases were more than twice those of non-poisoned incubations after 21 days. All other species showed significantly higher releases of TP after 1 day in the presence of poison (Table 3, Figure 2).

From Day 1 to Day 21, total release of TP by *S. alterniflora* blades increased, but the increases were most noticeable in the non-poisoned incubations, which more than doubled (Figure 2). In fact, after 21 days, poisoned and non-poisoned releases of TP by *S. alterniflora* were not significantly different (Table 3). After 21 days of incubation, all three Everglades species showed a significantly greater release of TP with poison (Table 3; Figure 2).

Release of TN could only be discerned in the non-poisoned incubations, as the NaN_3 poison interfered with our ability to detect significant TN change. Non-poisoned releases of TN from *Eleocharis spp.* culms were significantly greater than those by *R. mangle* and *C.*

jamaicense, but neither group could be distinguished from TN releases by *S. alterniflora*, the species we assumed to be limited by N in its natural environment (ANOVA; $p < 0.005$; Figure 3). Incubations containing *S. alterniflora* leaf material showed little net change in TN levels from Day 1 to Day 21, while the three Everglades species yielded more TN after 21 Days and showed similar trends in disparity between Day 1 and Day 21 (Figure 3)

Discussion

As expected, we saw much lower release of phosphorus from leaves of species collected in P-limited wetlands (Everglades National Park) compared to a N-limited wetland (North Inlet, SC). In fact, we estimate that the flux of P from leaching litter to a 1-m² patch of N-limited *S. alterniflora* ecosystem was up to two orders of magnitude greater than for the P-limited Everglades species (Table 4). Further, these releases were substantially lower in the presence of biological processes that appeared to prevent the release of that P to the water column.

Although we did not measure the P content of the leaf tissue (or the biofilm layer that developed on it), we assumed that the difference between TP release in poisoned and non-poisoned incubations was the result of microbes on the surface of the leaf mobilizing leached P. Our measurements of water column TOC, TN, and TP reflected the initial concentrations of these constituents plus the contribution of leached material and microbes suspended in the water column. Since we considered ‘total’ fractions in our analyses, declines in water column [TOC], [TN], and [TP] or differences between poisoned and non-poisoned incubations at a given time interval were assumed to be the result of respiration losses (C and N) or incorporation into microbial biomass on the leaf surface (C, N, and P).

2 The disparity between poisoned and non-poisoned treatments in our study was greatest in
3 the two freshwater Everglades macrophytes (*C. jamaicensis* and *Eleocharis spp.*) and in *S.*
4 *alterniflora*. Less TP leached to the water with live microbial communities, suggesting that a
5 significant portion of P leached from the plants in this study was labile and available to
6 microbes. It is likely that the leached P boosted microbial activity in these treatments, as P not
7 only limits primary production, but also microbial processes in this wetland (Davis, 1989,
8 Amador & Jones, 1993).

9 The same may be true for *S. alterniflora*. Although N limits primary productivity in
10 North Inlet (SC) salt marshes, Sundareshwar et al. (2003) showed that P limits soil microbial
11 respiration. This appeared to be the case early in our incubations when epiphytic microbial
12 communities on dead leaf material reduced P leaching into the water column by half (Figure 2).
13 However, as time progressed, molar ratios of TN:TP in *S. alterniflora* incubations suggested a
14 precipitous decline in N availability relative to P—with TN:TP averaging 11 in source water and
15 less than 2 after 21 days in poisoned incubations (Figure 4). This trend was supported by TP and
16 TN releases by *S. alterniflora* after 21 days.

17 That there was no difference between poisoned and non-poisoned release of TP after 21
18 days suggests that there was little demand for leached TP by epiphytic microbes at that time.
19 Further, no difference in the total amount of TN released by *S. alterniflora* in non-poisoned
20 incubations after 1 day versus after 21 days suggests that TN releases were diminished by
21 epiphytic microbes sometime between Day 1 and 21. Based on the change in molar TN:TP over
22 the duration of these incubations, it is likely that N became limiting to the microbial community
23 shortly after the first day of this set of incubations—when TN:TP averaged approximately 3
24 (Figure 4; Figure 6). All Everglades species showed a trend in water column TN:TP similar to *S.*

alterniflora, but 21-day ratios were still over an order of magnitude higher, indicating P-
2 limitation throughout the duration of the leaf incubations (Figure 4).

Dwarfed *Rhizophora mangle* in the southern Everglades is widely thought to be limited
4 by P availability (Koch & Snedaker, 1997; Feller et al., 1999). However, it may be that the lack
of a biological effect on TP release after 1 day was the result of water column microbes in this
6 wetland being limited by the availability of labile organic carbon. We observed a two-fold
reduction in TOC released from *R. mangle* leaves after one day when biological activity was
8 present (Figure 1). This corresponded with the lack of a 1-day difference in TP leaching
between poisoned and non-poisoned incubations. After the large, 1-day pulse of TOC from these
10 leaves, biological activity likely shifted back to P-limitation, as evidenced by the significant
difference between poisoned and non-poisoned releases of TP after 21 days and the lack of a
12 significant difference between poisoned and non-poisoned releases of TOC after this same period
of time had elapsed (Figures 1 and 2).

14 Dead *R. mangle* leaves represent a potentially large source of leachable OC to the
surrounding ecosystem (Table 4), supporting the observation that mangroves are a significant
16 source of DOM to Everglades estuarine ecosystems (Jaffé et al., 2004). Everglades mangrove
ecosystems have higher surface water DOC concentrations, and more of this DOM is potentially
18 labile compared to upstream oligotrophic freshwater wetlands (Maie et al., 2005). However,
freshwater oligotrophic wetlands of the Everglades can also be a net source of potentially labile
20 carbohydrates and proteinaceous material (Lu et al., 2003).

The different Everglades species exhibited a range of abiotic P leaching losses. The
22 maximum amount of TP leached from *Eleocharis* spp. in the first day in the poisoned bottles
whereas *S. alterniflora*, *C. jamaicense*, and *R. mangle* leached TP at progressively slower rates

through Day 21. The two freshwater Everglades species showed interesting contrasts in abiotic P leaching. Although both *C. jamaicense* and *Eleocharis* spp. leached similar amounts of TP after 1 day in the poisoned bottles, after 21 days *C. jamaicense* leached twice as much TP compared to *Eleocharis* spp., but this was a statistically insignificant difference. The greater inherent leachability of the P remaining in *C. jamaicense* litter after re-adsorption compared to *Eleocharis* spp. is surprising given the lower concentration of P in dead *C. jamaicense* leaves than *Eleocharis* spp. culms (Rubio & Childers, in review). In the end, the microbially mediated release of TP from non-poisoned *C. jamaicense* was nearly identical to non-poisoned *Eleocharis* spp. culms and *R. mangle* leaves.

Leaching of nutrients from *C. jamaicense* litter results in larger ecosystem fluxes despite similar mass-specific leaching rates from litter as *Eleocharis* spp. The annual net primary productivity of *C. jamaicense* is much greater than *Eleocharis* spp. in the Everglades, translating into a difference in litter production between these two herbaceous species (Daoust & Childers, 2004; Childers et al., in review; Table 4). This results in a larger flux of P from soil to plant litter in the water column via translocation in *C. jamaicense* marsh ($7.49 \text{ mmol P m}^{-2} \text{ yr}^{-1}$) compared to *Eleocharis* spp. sloughs ($1.10 \text{ mmol P m}^{-2} \text{ yr}^{-1}$; Noe & Childers, in review). The higher litter production of *C. jamaicense* compared to *Eleocharis* spp. also results in much larger fluxes of P, N, and C from this litter to the water column via leaching (Table 4). Using the non-poisoned leaching data from this study, we estimate the annual P flux from litter to the water column by leaching to be 0.69 and $0.10 \text{ mmol P m}^{-2} \text{ yr}^{-1}$ for *C. jamaicense* and *Eleocharis* spp., respectively. Thus, 9% of total litter P flux is leached into the water column for both species.

Phosphorus uptake from fresh litter by epiphytic microbes can be estimated by the difference in leaching fluxes between poisoned and non-poisoned treatments, equal to 1.55 and

0.07 mmol P m⁻² yr⁻¹ for *C. jamaicense* and *Eleocharis* spp., respectively, or 21% and 6% of total
2 litter P flux, respectively. This much larger flux of P to epiphytic microbes on *C. jamaicense*
compared to *Eleocharis* spp. could have large feedbacks on ecosystem P cycling and transport.
4 Phosphorus in microbial biomass is more labile and likely has faster turnover rates than the
refractory P remaining in plant litter, and could be more easily transported downstream through
6 long-term nutrient spiraling processes. The larger sum of P leaching fluxes from litter to both
the water column and epiphytic microbes in *C. jamaicense* marsh (2.23 mmol P m⁻² yr⁻¹)
8 compared to *Eleocharis* spp. sloughs (0.16 mmol P m⁻² yr⁻¹) represents a greater recycling of P
from *C. jamaicense* plants to other ecosystem components in the soil and water column of this
10 oligotrophic, P-limited ecosystem. Finally, the increase in the P content of decomposing litter in
the Everglades (Davis, 1991; Qualls & Richardson, 2000) could be explained, in part, by
12 microbial mobilization of P leaching from litter and not solely uptake of water column P.

14 **Conclusions**

Abiotic leaching accounted for the greatest loss of C, N, and P from leaves of the four
16 species we considered. As expected, TP release from P-limited Everglades plant species was
much lower than TP release by an N-limited North Inlet (SC) salt marsh plant. The presence of
18 microbial activity diminished the observed leaching yield, as normalized releases of C and P in
non-poisoned incubations were usually significantly lower than in the poisoned incubations.
20 This suggests that that biological degradation of leaves in wetland settings may in fact be
stimulated by the physical process of leaching that results in a rapid, labile energy source or
22 limiting elements such as phosphorus or nitrogen for microbial decomposers. Leaching of

nutrients from plant litter to the water column can represent a large flux of nutrients in
2 oligotrophic ecosystems.

Long-term Ecological Research in the Florida Coastal Everglades seeks to understand the
4 role of wetland hydrology in driving materials exchange (carbon, nitrogen, phosphorus,
suspended material, etc.) and productivity patterns at the land-sea interface. Our results suggest
6 that early leaf litter decay (i.e., leaching and microbial colonization) can contribute to local
regulation of surface water quality. However, analogous studies should be conducted in the field
8 to ascertain the actual contribution of early macrophyte decay to spatial and temporal patterns in
wetland ecosystem function. Further, studies such as these should also consider gradients of
10 ecosystem fertility and trophic status to fully understand the interactions between biological and
physical processes in the early decay of wetland plant tissue.

12 **Acknowledgements**

14 We thank Damon Rondeau and the Southeast Environmental Research Center for analytical
support. Jenny Davis, Luz Romero, and Alejandro Gaviria provided much-needed field and lab
16 support in completing these experiments. This work was supported in part by a 2002 Visiting
Scientist award to SED from the Belle Baruch Institute at the University of South Carolina and is
18 SERC contribution # XXX. This material is based upon work supported by the National Science
Foundation to the Florida Coastal Everglades LTER Program (Grant No. 9910514), the
20 Everglades Priority Ecosystem Science Initiative of the U.S. Geological Survey, and the National
Research Program of the U.S. Geological Survey.

List of References

2

Amador, J.A. & R.D. Jones. 1993. Nutrient limitations on microbial respiration in peat soils
4 with different total phosphorus content. *Soil Biology and Biochemistry*. 25(6): 793-801.

6

Aerts, R. 1996. Nutrient resorption from senescing leaves of perennials: are there general
patterns? *Journal of Ecology*. 84:597-608.

8

Benner, R., E.R. Peele, & R.E. Hodson. 1986. Microbial utilization of dissolved
10 organic matter from leaves of the red mangrove, *Rhizophora mangle*, in the Fresh Creek Estuary,
Bahamas. *Estuarine, Coastal and Shelf Science*. 23:607-619.

12

Childers, D.L., D. Iwaniec, D. Rondeau, G. Rubio, E. Verdon, & C. Madden. In review. Primary
14 productivity in Everglades marshes demonstrates the sensitivity of oligotrophic ecosystems to
environmental drivers. *Aquatic Botany*.

16

Daoust, R.J. & D.L. Childers, 2004. Ecological effects of low-level phosphorus additions on two
18 plant communities in a neotropical freshwater wetland ecosystem. *Oecologia*. 141:672-686.

20

Davis, S.E., C. Coronado-Molina, D.L. Childers, & J.W. Day, Jr. 2003. Temporally dependent
C, N, and P dynamics associated with the decay of *Rhizophora mangle* L. leaf litter in

22

oligotrophic mangrove wetlands of the southern Everglades. *Aquatic Botany*. 75:199-215.

- 2 Davis, S. M. 1989. Sawgrass and cattail production in relation to nutrient supply in the
3 Everglades. In R. R. S. and J. W. Gibbons (Eds.), *Freshwater Wetlands and Wildlife*. pp. 325-
4 341. U. S. Department of Energy. Charleston, S. C.
- 6 Davis, S.M. 1991. Growth, decomposition, and nutrient retention in *Cladium jamaicense* Crantz
7 and *Typha domingensis* Pers. in the Florida Everglades. *Aquatic Botany* 40: 203-224.
- 8 Feller, I. C., D. Whigham, J. O'Neill, & K. McKee. 1999. Effects of nutrient enrichment on
9 within-stand cycling in a mangrove forest. *Ecology*. 80(7):2193-2205.
- 10
- 11 Gosselink, J.G. & C.J. Kirby. 1974. Decomposition of salt marsh grass, *Spartina alterniflora*
12 Loisel. *Limnology and Oceanography*. 19(5):825-832.
- 14 Gosz, J.R., G.E. Likens, & F.H. Borman. 1973. Nutrient release from decomposing leaf and
15 branch litter in the Hubbard Brook Forest, New Hampshire. *Ecological Monographs*. 43:173-
16 191.
- 18 Ibrahima, A., R. Joffre, & D. Gillon. 1995. Changes in leaf litter during the initial leaching
19 phase: An experiment on the leaf litter of Mediterranean species. *Soil Biology and Biochemistry*.
20 27(7):931-939.

- 1 Jaffé, R., J.N. Boyer, X. Lu, N. Maie, C. Yang, N.M. Scully, & S. Mock. 2004. Source
2 characterization of dissolved organic matter in a subtropical mangrove-dominated estuary by
fluorescence analysis. *Marine Chemistry*. 84:195-210.
- 4
- Killingbeck, K. T. 1996. Nutrients in senesced leaves: keys to the search for potential resorption
6 and resorption proficiency. *Ecology*. 77(6):1716-1727.
- 8 Koch, M. S. & S. C. Snedaker. 1997. Factors influencing *Rhizophora mangle* L. seedling
development in Everglades carbonate soils. *Aquatic Botany*. 59:87-98.
- 10
- Lu, X., D.L. Childers, J.V. Hanna, N. Maie, & R. Jaffé. 2003. Molecular characterization of
12 dissolved organic matter in freshwater wetlands of the Florida Everglades. *Water Research*.
37:2599-2606.
- 14
- Maie, N., C. Yang, T. Miyoshi, K. Parish, & R. Jaffe. 2005. Chemical characteristics of
16 dissolved organic matter in an oligotrophic subtropical wetland/estuarine ecosystem. *Limnology
and Oceanography*. 50:23-35.
- 18
- Morris, J.T. & B. Haskin. 1990. A 5-yr record of aerial primary production and stand
20 characteristics of *Spartina alterniflora*. *Ecology* 71:2209-2217.
- 22 Noe, G.B., D.L. Childers, & R.D. Jones. 2001. Phosphorus biogeochemistry and the impact of
phosphorus enrichment: Why is the Everglades so unique? *Ecosystems*. 4:603-624.

- 2 Noe, G.B. & D.L. Childers. In review. Phosphorus budgets in Everglades wetland ecosystems:
The effects of nutrient enrichment. *Ecosystems*.
- 4
- 6 Qualls, R.G. & C.J. Richardson. 2000. P enrichment affects litter decomposition,
immobilization, and soil microbial P in wetland mesocosms. *Soil Science Society of America
Journal*. 64:799-808.
- 8
- 10 Richardson, C.J., G. Ferrell, & P. Vaithiyathan. 1999. Nutrient effects on stand structure,
resorption efficiency, and secondary compounds in Everglades sawgrass. *Ecology*. 80(7):2182-
2192.
- 12
- 14 Rivera-Monroy, V.H., S. Ewe, R.R. Twilley, C. Coronado-Molina, E. Castaneda, E., & T.J.
Grahl. In preparation. Scrub mangrove forest growth patterns and above-belowground
productivity in Taylor River, Everglades National Park, Florida, USA.
- 16
- 18 Romero, L.M., T.J. Smith, & J.W. Fourqurean. 2005. Changes in mass and nutrient content of
wood during decomposition in a south Florida mangrove forest. *Journal of Ecology*. 93:618-
631.
- 20
- 22 Rubio, G.A. & D.L. Childers. In Review. Decomposition of *Cladium jamaicense*, *Eleocharis*
sp., and *Juncus roemerianus* in the estuarine ecotones of the Florida Everglades. *Estuaries*.

- 2 Solorzano, L, & J, H, Sharp. 1980. Determination of total dissolved phosphorus and particulate
phosphorus in natural waters. *Limnology and Oceanography*. 25(4):754-758.
- 4 Sundareshwar, P.V., J.T. Morris, E.K. Koepfler, & B. Fornwalt. 2003. Phosphorus limitation of
coastal ecosystem processes. *Science*. 299:563-565.
- 6
- 8 Swift, M.J., O.W. Heal, & J.M. Anderson. 1979. *Decomposition in Terrestrial Ecosystems*.
Blackwell Scientific, Oxford, UK.
- 10 Taylor, B. R. & F. Bärlocher. 1996. Variable effects of air-drying on leaching losses from tree
leaf litter. *Hydrobiologia*. 325:173-182.
- 12
- 14 Teal, J. M. 1962. Energy flow in the salt marsh ecosystem of Georgia. *Ecology*. 43:614-624.
- 16 Tukey, H. B. 1970. The leaching of substances from plants. *Annual Review of Plant Physiology*.
21:305-324.
- 18 Webster, J.R. & E.F. Benfield. 1986. Vascular plant breakdown in freshwater ecosystems.
Annual Review of Ecology and Systematics 17: 567-594

2 Table 1: Physical location, description of collection sites, the ratio of oven-dried mass to air-dried mass (\pm stdev) for each species,
 and the date (month/year) of each collection and experiment. Site identifiers in parentheses refer to exact locations of FCE-LTER
 4 and North Inlet NERR sampling stations.

Site	Lat/Long (decimal degrees)	Wetland type	Fertility / Nutrient status	Species collected	Oven:Air mass (stdev)	Collection Date
SE Everglades, FL (TS/Ph 4)	25.315 N / -80.522 W	FW slough	oligotrophic, P- limited	<i>Cladium</i> <i>jamaicense</i>	0.879 (\pm 0.007)	Mar. 2001
SE Everglades, FL (TS/Ph 5)	25.295 N / -80.520 W	FW marsh	oligotrophic, P- limited	<i>Eleocharis</i> <i>spp.</i>	0.795 (\pm 0.006)	Apr. 2001
Taylor River, FL (TS/Ph 7b)	25.197 N / -80.642 W	mangrove	oligotrophic, P- limited	<i>Rhizophora</i> <i>mangle</i>	0.354 (\pm 0.019)	Aug. 2000
Clambank Creek, SC (North Inlet, CB)	33.334 N / -79.193 W	salt marsh	oligotrophic, N- limited	<i>Spartina</i> <i>alterniflora</i>	0.842 (\pm 0.004)	Jul. 2002

6

Table 2: Initial concentrations of salinity, TOC, TN, and TP in water sources used for each

2 leaching experiment.

Experiment (species)	Salinity (‰)	TOC (mM)	TN (μM)	TP (μM)
<i>C. jamaicense</i>	0	1.16	66.95	0.33
<i>Eleocharis</i> spp.	0	2.02	134.49	0.17
<i>R. mangle</i>	0	1.01	36.38	0.12
<i>S. alterniflora</i>	15	0.22	17.24	1.59

4

Table 3: *P*-values from one-factor ANOVAs testing the effect of poison on the release of TOC

2 and TP from each species after 1 day and after 21 days of incubation. In all significant tests ($P <$
 0.05), releases from leaves in poisoned bottles were greater than releases in non-poisoned bottles.
 4 ‘N.S.’ indicates no significant difference in the release of a given constituent between poisoned
 and non-poisoned bottles.

6

Species	1 Day (TOC)	21 Days (TOC)	Day 1 (TP)	21 Days (TP)
<i>R. mangle</i>	0.0060	N.S.	N.S.	0.0375
<i>C. jamaicense</i>	0.0052	0.0110	0.0129	0.0483
<i>Eleocharis spp.</i>	0.0028	0.0014	0.0117	0.0086
<i>S. alterniflora</i>	0.0002	0.0003	0.0010	N.S.

8

Table 4: Fluxes of nutrients in a representative 1-m² patch of wetland (mol m⁻² yr⁻¹) associated with the leaching of fresh detritus from different plant species, estimated from the product of mean leaching fluxes (mol g dw⁻¹) in this study and litterfall production values (g m⁻² yr⁻¹) from the literature. Leaching fluxes to water column are derived from the non-poisoned treatments, leaching fluxes to epiphytic microbes are derived as the difference between poisoned and non-poisoned treatments. Nitrogen fluxes were not measured in poisoned treatments. ^a Noe and Childers (in review); ^b Rivera-Monroy et al. (in preparation); ^c Morris and Haskin (1990).

Flux	<i>C. jamaicense</i>	<i>Eleocharis</i> spp.	<i>R. mangle</i>	<i>S. alterniflora</i>
Litterfall production (g dw m ⁻² yr ⁻¹)	1,789 ^a	281 ^a	120 ^b	635 ^c
Leaching flux to water column (mmol m ⁻² yr ⁻¹)				
TOC	870	90	4,500	610
P	0.69	0.10	0.05	6.93
N	25	4.8	1.4	7.0
Leaching flux to epiphytic microbes (mmol m ⁻² yr ⁻¹)				
TOC	1,400	330	1,300	1,300
P	1.55	0.07	0.08	0.62

8

List of Figures

2 Figure 1: Releases of total organic carbon (TOC) from each of four species of leaf material with
poison (gray bars) and without poison (white bars) incubated for 1 day (top graph) and for 21
4 days (bottom graph). Error bars represent standard deviations from three replicates. Axis break
for bottom graph reveals the disparity in TOC release between *R. mangle* and three macrophyte
6 species after 21 days of immersion in water.



8 Figure 2: Releases of total phosphorus (TP) from each of four species of leaf material with
poison (gray bars) and without poison (white bars) incubated for 1 day (top graph) and for 21
10 days (bottom graph). Error bars represent standard deviations from three replicates. Separate y-
axis for *S. alterniflora* is intended to show the magnitude of TP released from a N-limited plant
12 versus that of three P-limited, Everglades plant species after 1 and 21 days of immersion in
water.


14


Figure 3: Releases of total nitrogen (TN) from each of four species of leaf material immersed in
16 water without poison for 1 day (white bars) and 21 days (shaded bars). Error bars represent
standard deviations from three replicates.

18

Figure 4: Scatter plot showing how TN:TP molar ratios in incubation water changed through
20 time as a result of leaching and microbial activity associated with the early decay of all four
macrophyte species. Data are from non-poisoned treatments only and error bars represent
22 standard deviation of three replicates collected at each sampling interval.







2

