


2015 ANNUAL REPORT

OF THE WATER QUALITY MONITORING PROJECT

FOR THE WATER QUALITY PROTECTION PROGRAM

OF THE FLORIDA KEYS NATIONAL MARINE SANCTUARY



Principal Investigators:
Henry O. Briceño & Joseph N. Boyer¹

Southeast Environmental Research Center
OE-148, Florida International University
Miami, FL 33199. <http://serc.fiu.edu/wqmnetwork/>

¹Center for the Environment, Plymouth State University,
MSC 63, 17 High St., Plymouth, NH 03264

US EPA Agreement #X7 00D02412-1

This is Technical Report #799T of the Southeast Environmental Research Center,
Florida International University

June 2016

This page is intentionally left blank

2015 ANNUAL REPORT OF THE WATER QUALITY MONITORING PROJECT FOR THE WATER QUALITY PROTECTION PROGRAM OF THE FLORIDA KEYS NATIONAL MARINE SANCTUARY

Henry O. Briceño & Joseph N. Boyer

Funded by the Environmental Protection Agency (X7 00D02412-1)

EXECUTIVE SUMMARY

This report serves as a summary of our efforts to date in the execution of the Water Quality Monitoring Project for the FKNMS as part of the Water Quality Protection Program. The period of record for this report is Apr. 1995 – Dec. 2015 and includes data from 82 quarterly sampling events within the FKNMS (20.5 years). This annual report reflects funding cutbacks in 2012 resulting in reduction of spatial sampling from 155 to 112 sites.

Field parameters measured at each station (surface and bottom at most sites) include salinity (practical salinity scale), temperature ($^{\circ}\text{C}$), dissolved oxygen (DO, mg l^{-1}), turbidity (NTU), relative fluorescence, and light attenuation (K_d , m^{-1}). Water quality variables include the dissolved nutrients nitrate (NO_3^-), nitrite (NO_2^-), ammonium (NH_4^+), and soluble reactive phosphorus (SRP). Total unfiltered concentrations include those of nitrogen (TN), organic carbon (TOC), phosphorus (TP), silicate (SiO_2) and chlorophyll a (CHLA, $\mu\text{g l}^{-1}$).

The EPA developed Strategic Targets for the Water Quality Monitoring Project (SP-47) which state that beginning in 2008 through 2015, they shall annually maintain the overall water quality of the near shore and coastal waters of the FKNMS according to 2005 baseline. For reef sites, chlorophyll a should be less than or equal to $0.35 \mu\text{g l}^{-1}$ and the vertical attenuation coefficient for downward irradiance (K_d , i.e., light attenuation) should be less than or equal to 0.20 m^{-1} . For all monitoring sites in FKNMS, dissolved inorganic nitrogen should be less than or equal to $0.75 \mu\text{M}$ (0.010 ppm) and total phosphorus should be less than or equal to $0.25 \mu\text{M}$ (0.0077 ppm). Table 1 shows the number of sites and percentage of total sites exceeding these Strategic Targets for 2015.

We must recognize that the reduction of sampling sites in western FKNMS (less human-impacted sites) and the increase in inshore sites (heavily human-impacted sites) introduces a bias to the dataset which results in a reporting problem, perhaps requiring a revision of SP-47 to correct this deviation. To avoid such complications, we have not included the recently added locations (#500 to #509) in the calculation of compliances.

Table 1: EPA WQPP WQ Targets derived from 1995-2005 Baseline

For reef stations, chlorophyll less than or equal to 0.35 micrograms liter⁻¹ (ug l⁻¹) and vertical attenuation coefficient for downward irradiance (K_d , i.e., light attenuation) less than or equal to 0.20 per meter; for all stations in the FKNMS, dissolved inorganic nitrogen less than or equal to 0.75 micromolar and total phosphorus less than or equal to 0.25 micromolar; water quality within these limits is considered essential to promote coral growth and overall health. The “number of samples” exceeding these targets is tracked and reported annually. Values in **green** are those years with % compliance greater than 1995-2005 **baseline**. Values in **yellow** are those years with % compliance less than 1995-2005 **baseline**.

EPA WQPP Water Quality Targets				
	REEF Stations		All Stations (excluding SHORE sites)	
Year	CHLA ≤ 0.35 µg l ⁻¹	K_d ≤ 0.20 m ⁻¹	DIN ≤ 0.75 µM	TP ≤ 0.25 µM
			(0.010 ppm)	(0.0077 ppm)
1995-05	1778 of 2367 (75.1%)	1042 of 1597 (65.2%)	7826 of 10254 (76.3%)	7810 of 10267 (76.1%)
2006	196 of 225 (87.1%)	199 of 225 (88.4%)	432 of 990 (43.6%)	316 of 995 (31.8%)
2007	198 of 226 (87.6%)	202 of 222 (91.0%)	549 of 993 (55.3%)	635 of 972 (65.3%)
2008	177 of 228 (77.6%)	181 of 218 (83.0%)	836 of 1,000 (83.6%)	697 of 1,004 (69.4%)
2009	208 of 228 (91.2%)	189 of 219 (86.3%)	858 of 1,003 (85.5%)	869 of 1,004 (86.6%)
2010	170 of 227 (74.9%)	176 of 206 (85.4%)	843 of 1000 (84.3%)	738 of 1,003 (73.6%)
2011	146 of 215 (67.9%)	156 of 213 (73.2%)	813 of 1012 (80.3 %)	911 of 1013 (89.9 %)
2012	142 of 168 (84.5%)	135 of 168 (80.4%)	489 of 683 (71.6 %)	634 of 684 (92.7 %)
2013	148 of 172 (86.0%)	150 of 172 (87.2%)	496 of 688 (72.1 %)	603 of 688 (87.6 %)
2014	141 of 172 (82.0%)	133 of 172 (77.3%)	426 of 690 (61.7%)	540 of 690 (78.3%)
2015	122 of 172 (70.9%)	135 of 172 (78.5%)	487 of 688 (70.8%)	613 of 688 (89.1%)

Important Results Realized from this Monitoring Project

1. Land-based Influence on Water Quality

We documented elevated nutrient concentrations (DIN, TN, TP, and SiO₂) in waters close to shore along the Keys, and their corresponding responses from the system, such as higher phytoplankton biomass (CHLA), turbidity, as well as lower salinity and DO in the water column. These changes, associated to human impact, have become even more obvious by the addition of 10 stations located close to shore, (SHORE, within the 500 m Halo; FDEP 2011) sampled since Nov 2011 (Fig i).




Figure i. Nutrient and response changes along transect from shore sites to reef-track

This trend, especially for DIN was evident from our first sampling event in 1995 and was not observed in a comparison transect from the Tortugas (no human impact). This pattern suggests a land-based, freshwater nutrient source. In summary, this type of distribution would imply a relatively nutrient-rich land source which is diluted by low nutrient Atlantic Ocean waters.

2. Trend analysis – 20 years!

Surface DO has increased in all areas of the FKNMS (Fig. ii). Greatest increases in surface DO were generally observed on the Atlantic side of the Keys and in some inshore areas on the Bay side. Bottom DO trends showed a similar pattern as surface with more increased DO than surface sites (Fig. iii). Increased DO is beneficial for animal life.




Figure ii. Total change in DO of surface waters for 20 year period calculated from significant trends.




Figure iii. Total change in DO of bottom waters for 20 year period calculated from significant trends.

Water column turbidity (cloudiness) declined throughout the FKNMS during the 20 year period (this is good thing). There was no significant change in turbidity in bottom waters. The largest declines in turbidity occurred in western Florida Bay and west of the Marquesas (Fig iv).




Figure iv. Total change in Turbidity in surface waters for 20 year period calculated from significant trends.

Although K_d did not show significant trend, decreased turbidity affected the amount of light reaching the bottom (I_0 in %). I_0 increased at most reef sites throughout the Keys (Fig. v). More light on the bottom is beneficial to corals, seagrass, and algae. Interestingly, the Backcountry area of the lower Keys experienced increases in K_d which lead to corresponding decreases in I_0 .




Figure v. Total change in bottom I_0 for 20 year period calculated from significant trends.

No significant trends in TP were observed, however very small increases in SRP, up to 1.5 ppb over 20 years, were observed most surface waters (Fig. vi). Concentrations of SRP are generally an order of magnitude lower than TP and usually below kinetic uptake threshold of phytoplankton, meaning not all SRP is accessible.




Figure vi. Total change in SRP in surface waters for 20 year period calculated from significant trends.

The largest sustained monotonic trend has been the decline in surface TOC, especially in the Backcountry and the Marquesas (Fig. vii). This is part of a regional trend in TOC observed on the SW Shelf, Florida Bay, and the Everglades mangrove estuaries. This decline could be considered favorable given that TOC corresponds with CDOM (an important driver of water color and light penetration), but could also be an indication of decreased upstream primary production.




Figure vii. Total change in TOC in surface waters for 20 year period calculated from significant trends.

The large scale of this monitoring program has allowed us to assemble a much more holistic view of broad physical/chemical/biological interactions occurring over the South Florida hydroscape. This confirms that rather than thinking of water quality monitoring as being a static, non-scientific pursuit it should be viewed as a tool for answering management questions and developing new scientific hypotheses. We continue to maintain a website (<http://serc.fiu.edu/wqmnetwork/>) where data and reports from the FKNMS are integrated with other available programs.

Table of Contents

1.	Project Background	10
2.	Methods.....	13
2.1.	Field Sampling	13
2.2.	Laboratory Analysis	14
2.3.	Box and Whisker Plots.....	15
2.4.	Contour Maps.....	15
2.5.	Trend Analysis	15
3.	Results	16
3.1.	Overall Water Quality of the FKNMS in 2015	16
3.2.	Trend Analysis	20
4.	Overall Trends	24
4.1.	Strategic Targets	25
5.	2015 Condition Discussion	27
6.	Numeric Nutrient Criteria Development.....	51
6.1.	Acknowledgments.....	52
7.	References	53

1. Project Background

The Florida Keys are an archipelago of sub-tropical islands of Pleistocene origin which extend in a NE to SW direction from Miami to Key West and out to the Dry Tortugas (Fig. 1). In 1990, President Bush signed into law the Florida Keys National Sanctuary and Protection Act (HR5909) which designated a boundary encompassing >2,800 square nautical miles of islands, coastal waters, and coral reef tract as the Florida Keys National Marine Sanctuary (FKNMS). The Comprehensive Management Plan (NOAA 1995) required the FKNMS to have a Water Quality Protection Plan (WQPP) thereafter developed by EPA and the State of Florida (EPA 1995). The original agreement for the water quality monitoring component of the WQPP was subsequently awarded to the Southeast Environmental Research Program at Florida International University and the field sampling program began in March 1995.




Figure 1: Map of original FKNMS boundary including collapsed segment numbers and common names. Modified after Klein and Orlando (1994)

The waters of the FKNMS are characterized by complex water circulation patterns over both spatial and temporal scales with much of this variability due to seasonal influence in regional circulation regimes. The FKNMS is directly influenced by the Florida Current, the Gulf of Mexico Loop Current, inshore currents of the SW Florida Shelf (Shelf), discharge from the Everglades through the Shark River Slough, and by tidal exchange with both Florida Bay and Biscayne Bay (Lee et al. 1994, Lee et al. 2002).

Advection from these external sources has significant effects on the physical, chemical, and biological composition of waters within the FKNMS, as may internal nutrient loading and freshwater runoff from the Keys themselves (Boyer and Jones 2002). Water quality of the

FKNMS may be directly affected both by external nutrient transport and internal nutrient loading sources (Gibson et al. 2008). Therefore, the geographical extent of the FKNMS is one of political/regulatory definition and should not be thought of as an enclosed ecosystem.

A spatial framework for FKNMS water quality management was proposed on the basis of geographical variation of regional circulation patterns (Klein and Orlando, 1994). The final implementation plan (EPA 1995) partitioned the FKNMS into 9 sub-areas which was collapsed to 7 for routine sampling (Fig. 1). Station locations were developed using a stratified random design along onshore/offshore transects in sub-areas 5, 7, and 9 or within EMAP grid cells in sub-areas 1, 2, 4, and 6.

Sub-area 1 (Tortugas) includes the Dry Tortugas National Park (DTNP) and surrounding waters and is most influenced by the Loop Current and Dry Tortugas Gyre. Originally, there were no sampling sites located within the DTNP as it was outside the jurisdiction of NOAA. Upon request from the National Park Service, we initiated sampling at 5 sites within the DTNP boundary. Sampling in the Dry Tortugas was finally halted since 2011 due to budget constraints. Sub-area 2 (Marquesas) includes the Marquesas Keys and a shallow sandy area between the Marquesas and Tortugas called the Quicksands. Sub-area 4 (Backcountry) contains the shallow, hard-bottomed waters on the gulfside of the Lower Keys. Sub-areas 2 and 4 are both influenced by water moving south along the SW Shelf. Sub-area 6 can be considered as part of western Florida Bay. This area is referred to as the Sluiceway as it is strongly influenced by transport from Florida Bay, SW Shelf, and Shark River Slough (Smith, 1994). Sub-areas 5 (Lower Keys), 7 (Middle Keys), and 9 (Upper Keys) include the inshore, Hawk Channel, and reef tract of the Atlantic side of the Florida Keys. The Lower Keys are most influenced by cyclonic gyres spun off of the Florida Current, the Middle Keys by exchange with Florida Bay, while the Upper Keys are influenced by the Florida Current frontal eddies and to a certain extent by exchange with Biscayne Bay. All three oceanside segments are also influenced by wind and tidally driven lateral Hawk Channel transport (Pitts, 1997).

We have found that water quality monitoring programs composed of many sampling stations situated across a diverse hydroscape are often difficult to interpret due to the “can’t see the forest for the trees” problem (Boyer et al. 2000). At each site, the many measured variables are independently analyzed, individually graphed, and separately summarized in tables. This approach makes it difficult to see the larger, regional picture or to determine any associations among sites. In order to gain a better understanding of the spatial patterns of water quality of the FKNMS, we attempted to reduce the complicated data matrix into fewer elements which would provide robust estimates of condition and connection. To this end we developed an objective classification analysis procedure which grouped stations according to water quality similarity (Briceño et al. 2013, Fig. 2).




Figure 2: Map of FKNMS showing segments derived from Factor and Cluster Analysis of biogeochemical data: OFF=Offshore; MAR=Marquesas; BKS=Back Shelf; BKB= Back Bay; LK= Lower Keys; MK= Middle Keys; UK= Upper Keys

Although the original quarterly sampling of 155 stations was cut back to 112 (Fig. 3), it still provides a unique opportunity to explore the spatial component of water quality variability in the FKNMS, but eliminates the possibility of linking the Sanctuary's water quality to external sources of variability.




Figure 3. The SERC Water Quality Monitoring Network showing the distribution of fixed sampling stations within the FKNMS for 2015 sampling.

2. Methods

2.1. Field Sampling

The period of record of this study was from March 1995 to December 2015 which included 82 quarterly sampling events. For this year, field measurements and grab samples were collected from 112 fixed stations within the FKNMS boundary (Fig. 3). Depth profiles of temperature ($^{\circ}\text{C}$), salinity (practical salinity scale), dissolved oxygen (DO , mg l^{-1}), photosynthetically active radiation (PAR , $\mu\text{E m}^{-2} \text{s}^{-1}$), chlorophyll a specific fluorescence (FSU), turbidity (NTU), depth as measured by pressure transducer (m), and density (σ_t , in kg m^{-3}) were measured by CTD casts (Seabird SBE 19). The CTD was equipped with internal RAM and operated in stand-alone mode at a sampling rate of 0.5 sec. The vertical attenuation coefficient for downward irradiance (K_d , m^{-1}) was calculated at 0.5 m intervals from PAR and depth using the standard exponential equation (Kirk 1994) and averaged over the station depth. This was necessary due to periodic occurrence of optically distinct layers within the water column. During these events, K_d was reported for the upper layer. To determine the extent of stratification we calculated the difference between surface and bottom density as delta Sigma-t ($\Delta\sigma_t$, in kg m^{-3}), where positive values denoted greater density of bottom water relative to the surface. A $\Delta\sigma_t > 1$ is considered weakly stratified, while any instances > 2 is strongly stratified.

In the Backcountry area (Sub-area 4, Fig. 1) where it is too shallow to use a CTD, surface salinity and temperature were measured using a combination salinity-conductivity-temperature-DO probe (YSI 650 MDS display-datalogger with YSI 6600V2 sonde). DO was automatically corrected for salinity and temperature. PAR was measured every 0.5 m using a

Li-Cor LI-1400 DataLogger equipped with a 4π spherical sensor (LI-193SB). PAR data with depth was used to calculate K_d from in-air surface irradiance.

Water was collected from approximately 0.25 m below the surface and at approximately 1 m from the bottom with a Niskin bottle (General Oceanics) except in the Backcountry and Sluiceway where surface water was collected directly into sample bottles. Duplicate, unfiltered water samples were dispensed into 3x sample rinsed 120 ml HDPE bottles for analysis of total constituents. Duplicate water samples for dissolved nutrients were dispensed into 3x sample rinsed 150 ml syringes which were then filtered by hand through 25 mm glass fiber filters (Whatman GF/F) into 3x sample rinsed 60 ml HDPE bottles. The resulting wet filters, used for chlorophyll *a* (CHLA) analysis, were placed in 1.8 ml plastic centrifuge tubes to which 1.5 ml of 90% acetone/water was added (Strickland and Parsons 1972).

All samples were kept on ice in the dark during transport to the laboratory. During overnight stays in the Lower Keys sampling, filtrates and filters (not total samples) were frozen until further analysis.

2.2. Laboratory Analysis

Samples were analyzed for ammonium (NH_4^+), nitrate+nitrite (N+N), nitrite (NO_2^-), total nitrogen (TN), soluble reactive phosphorus (SRP), total phosphorus (TP), total organic carbon (TOC), silicate (SiO_2), chlorophyll *a* (CHLA, $\mu\text{g l}^{-1}$), and turbidity (NTU) using standard laboratory methods. Dissolved nutrients were defined using Whatman GF/F filters with a nominal pore size of 0.8 μm . A 60 ml sample was collected from a Niskin bottle using a syringe and filtered through a 25 mm Whatman GF/F filter. The filtrate was collected in a 60 ml high density polyethylene (HDPE) bottle and the filter stored in a vial with 90% acetone for extraction of CHLA. An additional 120 ml sample was collected directly from the Niskin bottle for analysis of TN, TP, and turbidity.

NH_4^+ was analyzed by the indophenol method (Koroleff 1983). NO_2^- was analyzed using the diazo method and N+N was measured as nitrite after cadmium reduction (Grassoff 1983a,b). The ascorbic acid/molybdate method was used to determine SRP (Murphy and Riley 1962). High temperature combustion and high temperature digestion were used to measure TN (Frankovich and Jones 1998; Walsh 1989) and TP (Solórzano and Sharp 1980), respectively. TOC was determined using the high temperature combustion method of Sugimura and Suzuki (1988). Silicate was measured using the heteropoly blue method (APHA 1995). Samples were analyzed for CHLA content by spectrofluorometry of acetone extracts (Yentsch and Menzel 1963). Protocols are presented in EPA (1993) and elsewhere as noted. All elemental ratios discussed were calculated on a molar basis. DO saturation in the water column (DO_{sat} as %) was calculated using the equations of Garcia and Gordon (1992). Some parameters were not measured directly but calculated by difference. Nitrate (NO_3^-) was calculated as $\text{N+N} - \text{NO}_2^-$; total dissolved inorganic nitrogen (DIN) as $\text{N+N} + \text{NH}_4^+$, and total organic nitrogen (TON) as $\text{TN} - \text{DIN}$. All variables are reported in ppm (mg l^{-1}) unless otherwise noted.

In accordance with EPA policy, the FKNMS water quality monitoring program adhered to existing rules and regulations governing QA and QC procedures as described in EPA guidance documents. The FIU-SERC Nutrient Laboratory maintained NELAP certification during this project.

2.3. Box and Whisker Plots

Typically, water quality data are skewed to the left (low concentrations and below detects) resulting in non-normal distributions. Therefore it is more appropriate to use the median as the measure of central tendency because the mean is inflated by high outliers (Christian et al. 1991). Data distributions of water quality variables are reported as box-and-whiskers plots. The box-and-whisker plot is a powerful statistic as it shows the median, range, the data distribution as well as serving as a graphical, nonparametric ANOVA. The center horizontal line of the box is the median of the data, the top and bottom of the box are the 25th and 75th percentiles (quartiles), and the ends of the whiskers are the 5th and 95th percentiles. The notch in the box is the 95% confidence interval of the median. When notches between boxes do not overlap, the medians are considered significantly different. Outliers (<5th and >95th percentiles) were excluded from the graphs to reduce visual compression. Differences in variables were also tested between groups using the Wilcoxon Ranked Sign test (comparable to a *t*-test) and among groups by the Kruskal-Wallis test (ANOVA) with significance set at $p < 0.05$.

2.4. Contour Maps

In an effort to elucidate the contribution of external factors to the water quality of the FKNMS and to visualize gradients in water quality over the region, we combined Keys and Shelf data into contour maps (ArcView, ESRI) of specific water quality variables until 2011, when monitoring of the Shelf was dropped. We used kriging as the geostatistical algorithm because it is designed to minimize the error variance while at the same time maintaining point pattern continuity (Isaaks & Srivastava, 1989). Kriging is a general method of statistical interpolation that can be applied within any discipline to sampled data from random fields that satisfy the appropriate mathematical assumptions. Kriging is a global approach which uses standard geostatistics to determine the "distance" of influence around each point and the "clustering" of similar samples sites (autocorrelation). Therefore, unlike the inverse distance procedure, kriging will not produce valleys in the contour between neighboring points of similar value.

2.5. Trend Analysis

Temporal trends (unit yr^{-1}) were quantified by simple linear regression. Trend graphs were drawn only for those variables for which 15% of individual station trends were significant ($p < 0.10$). In an effort to show trend impact, significant trends are reported as the total change in concentration over the period of record.

3. Results

3.1. Overall Water Quality of the FKNMS in 2015

Summary statistics for all water quality variables from calendar year 2015 sampling events are shown as number of samples (n), minimum, maximum, and median (Table 1). Overall, the region remains warm and euhaline with a median temperature of 26.4 °C and salinity of 36.3; dissolved oxygen saturation of the water column (DO_{sat}) was relatively high at 97.1%. On this coarse scale, the FKNMS exhibited very good water quality with median NO_3^- , NH_4^+ , TP, and SiO_2 concentrations of 0.0028, 0.0049, 0.0057, and 0.008 mg l^{-1} , respectively. NH_4^+ was the dominant DIN species in almost all of the samples (~70%). However, DIN comprised a small fraction (7.5%) of the TN pool (0.115 mg l^{-1}) with TON being the bulk (median 0.107 mg l^{-1}). SRP concentrations were low (median 0.0011 mg l^{-1}) and comprised 19% of the TP pool (0.0057 mg l^{-1}). CHLA concentrations were also low overall, 0.28 $\mu\text{g l}^{-1}$, but ranged from 0.07 to 8.76 $\mu\text{g l}^{-1}$. TOC was 1.46 mg l^{-1} ; a value higher than open ocean levels but consistent with coastal areas.

Median turbidity was higher than usual (0.31 NTU) compared to last year's median of 0.06 NTU. Interestingly, K_d was still low (0.225 m^{-1}), slightly up from last year. Overall, 33.7% of incident light (I_0) reached the bottom, which is down from last year's 38.1%. Molar ratios of N to P suggested a general P limitation of the water column (median TN:TP = 45.2) but this must be tempered by the fact that much of the TN may not be bioavailable.

Table 1. Summary statistics for each water quality variable in the FKNMS for the calendar year 2015. Data are summarized as number of samples (*n*), minimum value (*Min.*), maximum value (*Max.*), and Median.

Variable	Depth	<i>n</i>	Min.	Max.	Median
NO₃⁻ (mg l ⁻¹)	Surface	407	0.0001	0.0175	0.0028
	Bottom	279	0.0001	0.0163	0.0027
NO₂⁻ (mg l ⁻¹)	Surface	407	0.0001	0.0027	0.0004
	Bottom	279	0.0001	0.0020	0.0004
NH₄⁺ (mg l ⁻¹)	Surface	407	0.0004	0.0240	0.0049
	Bottom	279	0.0008	0.0273	0.0047
TN (mg l ⁻¹)	Surface	407	0.0081	0.3930	0.1154
	Bottom	279	0.0001	0.2576	0.0986
DIN (mg l ⁻¹)	Surface	407	0.0018	0.0341	0.0087
	Bottom	279	0.0016	0.0297	0.0085
TON (mg l ⁻¹)	Surface	407	0.0001	0.3727	0.1067
	Bottom	279	0.0001	0.2529	0.0899
TP (mg l ⁻¹)	Surface	407	0.0009	0.0241	0.0057
	Bottom	279	0.0012	0.0218	0.0054
SRP (mg l ⁻¹)	Surface	406	0.0001	0.0133	0.0011
	Bottom	279	0.0001	0.0077	0.0010
CHLA (µg l ⁻¹)	Surface	406	0.07	8.76	0.28
TOC (mg l ⁻¹)	Surface	407	0.78	4.93	1.46
	Bottom	279	0.75	2.96	1.29
SiO₂ (mg l ⁻¹)	Surface	407	0.0001	0.937	0.008
	Bottom	279	0.0001	0.487	0.005
Turbidity (NTU)	Surface	390	0.01	6.09	0.31
	Bottom	265	0.01	5.18	0.30
Salinity	Surface	401	32.22	39.38	36.31
	Bottom	401	32.22	39.36	36.41
Temp. (°C)	Surface	401	22.64	32.97	26.43
	Bottom	401	22.61	32.98	26.18
DO (mg l ⁻¹)	Surface	401	5.73	9.09	6.49
	Bottom	401	5.84	9.16	6.55
K_d		402	0.001	1.102	0.225
TN:TP	Surface	407	2.9	362.5	45.2
Si:DIN	Surface	407	0.0	47.8	0.5
DO Saturation (%)	Surface	401	85.1	135.2	97.1
	Bottom	401	88.2	136.1	97.5
I₀ (%)	Bottom	400	1.3	99.1	33.7
Δσ_t		401	-0.028	2.118	0.017

The recently implemented stations close to shore were not used for this classification. In their short life-span they have displayed a common tendency to be nutrient-enriched and at the lower salinity extreme, as compared to the rest of the sites. Hence, we have grouped these ten stations as an additional class (SHORE), not for mapping purposes at this time, but for comparison and exploration of human impacts on water quality.

Most differences among segments were rather subtle, BKB, BKS and SHORE are the most nutrient-enriched segments while the UK and OFF were at the less-enriched extreme. The BKB zone was composed primarily of stations located inside and north of the Lower Keys and extending to the Sluiceway (Fig. 4). This class was highest in nutrients, especially TN, TON, TOC, SiO₂, TP, TOC and DIN, leading to high CHLA and turbidity. In the shallow BKB sites we expect that either nutrient transport from the SW Shelf and south Florida Bay and/or benthic flux of nutrients might be more important than anthropogenic loading. The BKB also had highest salinity and DO, relative to other regions. The BKS is located to the north of BKB and includes sites most influenced by water moving south from the SW Florida Shelf and exchange with BKB waters. It was highest in TP and relatively high in TN, TON, SRP, SiO₂, TOC, DO, turbidity and salinity.

The MAR zone was made up of sites between Key West and Rebecca Shoals. This is an area of relatively shallow water with complex circulation pattern which separates the SW Shelf from the Atlantic Ocean. The water quality of MAR is very low in TOC and relatively low in all N species and SiO₂, but displays relatively high TP and SRP, and the highest values and the largest range of variability in CHLA and turbidity, perhaps linked to shallow waters and sediment re-suspension.

There is a general nutrient gradient from higher levels at LK to MK to the UK, the less enriched one. Additionally, these three segments, closer to the islands, have higher nutrient levels than those offshore (OFF), underscoring the impact on water quality from the Keys and the strong control exerted by the Loop and Florida currents. The LK, MK and UK included the innermost sites of the Keys, which are shallow, closest to any possible anthropogenic nutrient sources, and typically more turbid than reef zones (OFF) from beach wave re-suspension. These sites were slightly elevated in DIN, TN, TON, SiO₂ and TOC relative to the OFF sites.

The OFF zone was made up of all Hawk Channel and reef tract sites of the mainland Keys and all sites west of Rebecca Shoal, including those in Dry Tortugas National Park. This zone had very low nutrients, TP, CHLA, and turbidity.




Figure 4. Box-and-whisker plots of surface samples showing median and distribution of water quality parameters as stratified by water quality cluster. Notches in the box that do not overlap with another are considered significantly different.

3.2. Trend Analysis

We must always keep in mind that trend analysis is limited to the window of observation; trends change with continued data collection. In addition, water quality in the Keys is largely externally-driven and may fluctuate according to climatic or disturbance events of longer periodicity. Trends may even reverse during a period of record. Examples of this are shown in Figures 5-7, where trends can be seen to be 1) monotonic, 2) episodically driven, and 3) reversing with change point.




Figure 5. Monotonic trend in TOC at Carysfort Reef.




Figure 6. Episodically (hurricane) driven pattern in NH_4^+ at The Elbow.




Figure 7. Reversing trend in DO at Carysfort Reef.

Least squares regressions for each water quality variable were calculated for the 20 year period of record. Only slopes having significant trends ($p < 0.10$) in ppm yr^{-1} , or as noted were reported; non-significant trends were coded as slope = 0. Some of the slopes are very small,

but to get an idea of total change over the period of record, the annual slopes were multiplied by number of years samples and plotted as contour maps of Change for 20 year period (Fig. 8-18). Only those variables with >20% of sampling sites having significant slopes were displayed.

Surface DO has increased in all areas of the FKNMS (Fig. 8). Greatest increases in surface DO were generally observed on the Atlantic side of the Keys and in some inshore areas on the Atlantic side. Bottom DO trends showed a similar pattern as surface with more increased DO than surface sites (Fig. 9). Increased DO is beneficial for animal life.




Figure 8. Total change in DO of surface waters for 20 year period calculated from significant trends.




Figure 9. Total change in DO of bottom waters for 20 year period calculated from significant trends.

Water column turbidity (cloudiness) declined throughout the FKNMS (a beneficial result) during the 20 year period (Fig 10). There was no significant change in turbidity in bottom waters. The largest declines in turbidity occurred in western Florida Bay and Marquesas.




Figure 10. Total change in surface turbidity for 20 year period calculated from significant trends.

Although K_d did not show significant trend, decreased turbidity increased the amount of light reaching the bottom (I_o in %). I_o increased mostly at reef sites throughout the Keys (Fig. 11). More light on the bottom is beneficial to corals, seagrass, and algae. Interestingly, the Backcountry area of the lower Keys experienced increases in K_d which lead to corresponding decreases in I_o and therefore less light on the bottom.




Figure 11. Total change in bottom I_o for 20 year period calculated from significant trends.

No significant trends in TP were observed, however very small increases in SRP, up to 1.5 ppb over 20 years, were observed most surface waters (Fig. 12). Concentrations of SRP are generally an order of magnitude lower than TP and usually below kinetic uptake threshold of phytoplankton, meaning not all SRP is accessible.




Figure 12. Total change in SRP in surface waters for 20 year period calculated from significant trends.

Clearly, there have been some changes in the FKNMS water quality over time, but the largest sustained monotonic trend has been the decline in surface TOC concentration. There were strong declines in surface TOC throughout the FKNMS, especially in the Backcountry and the Marquesas (Fig. 13). This is part of a regional trend in TOC observed on the SW Shelf, Florida Bay, and the mangrove estuaries draining the Everglades. This decline could be considered favorable given that TOC corresponds with CDOM (an important driver of water color and light penetration), but could also be an indication of decreased upstream primary production.




Figure 13. Total change in TOC in surface waters for 20 year period calculated from significant trends.

4. Overall Trends

Several important results have been realized from this monitoring project. First we documented elevated nutrient concentrations (DIN, TN, TP, and SiO₂) in waters close to shore along the Keys, and their corresponding responses from the system, such as higher phytoplankton biomass (CHLA), turbidity, as well as lower salinity and DO in the water column (Figure 14). These changes, associated to human impact, have become even more obvious by the addition of 10 stations (# 500 to #509) located very close to shore, sampled since Nov 2011 (SHORE).

Clearly, there have been some changes in the FKNMS water quality over time, and some sustained monotonic trends have been observed, however, we must always keep in mind that trend analysis is limited to the window of observation. Trends may change, or even reverse, with additional data collection. This brings up an important point that, when looking at what are perceived to be local trends, we find that they may occur across the whole region at more subtle levels. This spatial autocorrelation in water quality is an inherent property of highly interconnected systems such as coastal and estuarine ecosystems driven by similar hydrological and climate forcing. It is clear that trends observed inside the FKNMS are influenced by regional conditions outside the Sanctuary boundaries.




Figure 14: Nutrient and response changes along transect from close-to-shore sites to the reef-track

4.1. Strategic Targets

The EPA developed Strategic Targets for the Water Quality Monitoring Project which state that beginning in 2008, annually maintain the overall water quality of the near shore and coastal waters of the FKNMS according to 2005 baseline. For reef sites, chlorophyll *a* should be less than or equal to 0.2 micrograms/l and the vertical attenuation coefficient for downward irradiance (K_d , i.e., light attenuation) should be less than or equal to 0.13 per meter. For all monitoring sites in FKNMS, dissolved inorganic nitrogen should be less than or equal to 0.75 micromolar (0.010 milligrams/l) and total phosphorus should be less than or equal to 0.2 micromolar (0.0077 milligrams/l). Table 2 shows the number of sites and percentage of total sites exceeding these Strategic Targets for 2014.

Table 1: EPA WQPP WQ Targets derived from 1995-2005 Baseline

For reef stations, chlorophyll less than or equal to 0.35 micrograms liter⁻¹ (ug l⁻¹) and vertical attenuation coefficient for downward irradiance (K_d , i.e., light attenuation) less than or equal to 0.20 per meter; for all stations in the FKNMS, dissolved inorganic nitrogen less than or equal to 0.75 micromolar and total phosphorus less than or equal to 0.25 micromolar; water quality within these limits is considered essential to promote coral growth and overall health. The “number of samples” exceeding these targets is tracked and reported annually. Values in **green** are those years with % compliance greater than 1995-2005 **baseline**. Values in **yellow** are those years with % compliance less than 1995-2005 **baseline**.

EPA WQPP Water Quality Targets				
	REEF Stations		All Stations (excluding SHORE sites)	
Year	CHLA ≤ 0.35 µg l ⁻¹	K_d ≤ 0.20 m ⁻¹	DIN ≤ 0.75 µM	TP ≤ 0.25 µM
			(0.010 ppm)	(0.0077 ppm)
1995-05	1778 of 2367 (75.1%)	1042 of 1597 (65.2%)	7826 of 10254 (76.3%)	7810 of 10267 (76.1%)
2006	196 of 225 (87.1%)	199 of 225 (88.4%)	432 of 990 (43.6%)	316 of 995 (31.8%)
2007	198 of 226 (87.6%)	202 of 222 (91.0%)	549 of 993 (55.3%)	635 of 972 (65.3%)
2008	177 of 228 (77.6%)	181 of 218 (83.0%)	836 of 1,000 (83.6%)	697 of 1,004 (69.4%)
2009	208 of 228 (91.2%)	189 of 219 (86.3%)	858 of 1,003 (85.5%)	869 of 1,004 (86.6%)
2010	170 of 227 (74.9%)	176 of 206 (85.4%)	843 of 1000 (84.3%)	738 of 1,003 (73.6%)
2011	146 of 215 (67.9%)	156 of 213 (73.2%)	813 of 1012 (80.3 %)	911 of 1013 (89.9 %)
2012	142 of 168 (84.5%)	135 of 168 (80.4%)	489 of 683 (71.6 %)	634 of 684 (92.7 %)
2013	148 of 172 (86.0%)	150 of 172 (87.2%)	496 of 688 (72.1 %)	603 of 688 (87.6 %)
2014	141 of 172 (82.0%)	133 of 172 (77.3%)	426 of 690 (61.7%)	540 of 690 (78.3%)
2015	122 of 172 (70.9%)	135 of 172 (78.5%)	487 of 688 (70.8%)	613 of 688 (89.1%)

5. 2015 Condition Discussion

Water quality is a subjective measure of ecosystem well-being. Aside from the physical-chemical composition of the water there is also a human perceptual element which varies according to our intents for use (Kruczynski and McManus 2002). Distinguishing internal from external sources of nutrients in the FKNMS is a difficult task. The finer discrimination of internal sources into natural and anthropogenic inputs is even more difficult. Most of the important anthropogenic inputs are regulated and most likely controlled by management activities, however, recent studies have shown that nutrients from shallow sewage injection wells may be leaking into nearshore surface waters (Corbett et al. 1999; Shinn 1999a, 1999b; Paul et al. 1995, 1997; Reich et al. 2001; Briceño et al. 2015). Advective transport of nutrients through the FKNMS was not measured by the existing fixed sampling plan. However, nutrient distribution patterns may be compared to the regional circulation regimes in an effort to visualize the contribution of external sources and advective transport to internal water quality of the FKNMS (Boyer and Jones 2002).

Circulation in coastal South Florida is dominated by regional currents such as the Loop Current, Florida Current, and Tortugas Gyre and by local transport via Hawk Channel and along-shore Shelf movements (Klein and Orlando 1994). Regional currents may influence water quality over large areas by the advection of external surface water masses into and through the FKNMS (Lee et al. 1994, Lee et al. 2002) and by the intrusion of deep offshore ocean waters onto the reef tract as internal bores (Leichter et al. 1996). Local currents become more important in the mixing and transport of freshwater and nutrients from terrestrial sources (Smith 1994; Pitts 1997, Gibson et al. 2008).

Spatial patterns of salinity in coastal South Florida show these major sources of freshwater to have more than just local impacts (Fig. 15). In Biscayne Bay, freshwater is released through the canal system operated by the South Florida Water Management District; the impact may sometimes be seen to affect northern Key Largo by causing episodic depressions in salinity at alongshore sites. Freshwater entering NE Florida Bay via overland flow from Taylor Slough and C-111 basin mix in a SW direction. The extent of influence of freshwater from Florida Bay on alongshore salinity in the Keys is less than that of Biscayne Bay but it is more episodic. Transport of low salinity water from Florida Bay does not affect the Middle Keys sites enough to depress the median salinity in this region but is manifested as increased variability. The opposite also holds true; hypersaline waters from Florida Bay may be transported through the Sluiceway to inshore sites in the Middle Keys.

On the west coast, the large influence of the Shark River Slough, which drains the bulk of the Everglades and exits through the Whitewater Bay - Ten Thousand Islands mangrove complex, clearly impacts the Shelf waters. The mixing of Shelf waters with the Gulf of Mexico produces a salinity gradient in a SW direction which extends out to Key West. This freshwater source may sometimes affect the Backcountry because of its shallow nature but often follows a trajectory of entering western Florida Bay and exiting out through the channels in the Middle

Keys (Smith 1994). This net transport of lower salinity water from mainland to reef in open channels through the Keys is observed as an increase in the range and variability of salinity rather than as a large depression in salinity. All these forces have large influence on other water quality variables, especially DO (Fig. 16). Lowest DO concentrations tend to develop inside the Backcountry during warmest months.

In addition to surface currents there is evidence that internal tidal bores regularly impact the Key Largo reef tract (Leichter et al. 1996; Leichter and Miller 1999). Internal bores are episodes of higher density, deep water intrusion onto the shallower shelf or reef tract. Depending on their energy, internal tidal bores can promote stratification of the water column or cause complete vertical mixing as a breaking internal wave of sub-thermocline water.




Figure 15. Surface salinity distributions across the region during 2015.




Figure 16. Surface dissolved oxygen distributions across the region during 2015.

Visualization of spatial patterns of NO_3^- concentrations over South Florida waters provides an extended view of source gradients over the region (Fig. 17). The oceanside transects off the uninhabited Upper Keys (off Biscayne Bay) exhibited the lowest alongshore NO_3^- compared to the Middle and Lower Keys. A similar pattern was observed in a previous transect survey from these areas (Szmant and Forrester 1996). They also showed an inshore elevation of NO_3^- relative to Hawk Channel and the reef tract which is also demonstrated for DIN in our previous analysis (Fig. 4).

A distinct intensification of NO_3^- occurs in the Backcountry region. Part of this increase may be due to local sources of NO_3^- , i.e. septic systems and stormwater runoff around Big Pine Key (Lapointe and Clark 1992). However, there is another area, the Snipe Keys, that also exhibits high NO_3^- which is uninhabited by man, which rules out the premise of septic systems being the only source of NO_3^- in this area. It is important to note that the Backcountry area is very shallow (~0.5 m) and hydraulically isolated from the Shelf and Atlantic which results in its having a relatively long water residence time. Elevated NO_3^- concentrations may be partially due to simple evaporative concentration as is seen in locally elevated salinity values. Another possibility is a contribution of benthic N_2 fixation/nitrification in this very shallow area.

The elevated DIN concentrations in the Backcountry are not easily explained. We think that the high concentrations found there are due to a combination of anthropogenic loading, physical entrapment, and benthic N_2 fixation. The relative contribution of these potential sources is unknown. Lapointe and Matzie (1996) have shown that stormwater and septic systems are responsible for increased DIN loading in and around Big Pine Key. The effect of increased water residence time in DIN concentration is probably small. Salinities in this area were only 1-2 higher than local seawater which resulted in a concentration effect of only 5-6%.

Benthic N_2 fixation may potentially be very important in the N budget of the Backcountry. Measured rates of N_2 fixation in a *Thalassia* bed in Biscayne Bay, having very similar physical and chemical conditions, were $540 \mu\text{mol N m}^{-2} \text{d}^{-1}$ (Capone and Taylor 1980). Without the plant community N demand, one day of N_2 fixation has the potential to generate a water column concentration of $>0.014 \text{ ppm NH}_4^+$ (0.5 m deep). Much of this NH_4^+ is probably nitrified and may help account for the elevated NO_3^- concentrations observed in this area as well (Fig. 18). Clearly, N_2 fixation may be a significant component of the N budget in the Backcountry and that it may be exported as DIN to the FKNMS in general.




Figure 17. Surface nitrate distributions across the region during 2015.

Surface and bottom water concentrations are not always coincident. Interestingly, in many cases for 2015 and other years, we observe elevated NO_3^- in the bottom waters on the offshore reef tract (Fig. 18). We attribute this to regular “upwelling” (actually internal tidal bores) of deep water onto the reef tract (Leichter et al. 2003). This deep ocean water transport is a regular and persistent phenomenon which may deliver high nutrient waters to the offshore reef tract independent of any anthropogenic source.

During the Fall 2015 sampling event, an extensive water mass north of the Middle and Lower Keys exhibited relatively high NO_3^- concentrations. As we have no measurements outside of the FKNMS, we cannot speculate as to its source on the SW Shelf.




Figure 18. Bottom nitrate distributions across the region during 2015.

NH_4^+ concentrations were distributed in a similar manner as NO_3^- with highest levels occurring in the Backcountry (Fig. 19). NH_4^+ also showed additional similarities with NO_3^- in its spatial distribution, being lowest in the Upper Keys and highest inshore relative to offshore. This pattern of decline offshore implies an onshore N source which is diluted with distance from land by low nutrient Atlantic Ocean waters.

In many situations, independent water masses may be distinguished by difference in density (σ_t) between surface and bottom ($\Delta\sigma_t$, Fig. 20). Since density is driven more by salinity than temperature, we do not always observe differences in σ_t between surface and bottom during upwelling events. However, decreased temperature of bottom waters from intrusion of deeper oceanic waters is clearly an indicator of increased NO_3^- . These upwelling events also affect other nutrient species such as NH_4^+ , TP, and SRP in these bottom waters as well.




Figure 19. Surface ammonium distributions across the region during 2015.




Figure 20. Surface and bottom density differences ($\Delta\sigma_t$) across the region during 2015.

Spatial patterns in TP in South Florida coastal waters are strongly driven by the west coast sources (Boyer and Briceño 2007, 2011). A gradient in TP typically extends from the inshore waters of Whitewater Bay - Ten Thousand Islands mangrove complex out onto the Shelf and Tortugas. Gradients also extend from western Florida Bay to the Middle/Lower Keys. The spatial distribution of TP on the Shelf is driven by freshwater inputs from mangrove rivers and transport of Gulf of Mexico waters through the region. No significant evidence of a groundwater source exists (Corbett et al. 2000). Little can be concluded regarding TP distribution in the Sanctuary during 2015, except that the highest concentrations (between 0.005 and 0.010 mg l⁻¹ TP) preferentially occurred on the Bay side of the Keys, and were probably supplied by Shelf waters (Fig. 21). Also, in winter 2015, deeper offshore waters may have contributed TP to shallower localities in the Upper Keys.




Figure 21. Distributions of surface total phosphorus across the region during 2015.

Concentrations of TOC (Fig. 22) and TN (Fig. 23) are similar in pattern of distribution across the South Florida coastal hydroscape but do not always correspond. We believe this is due to differences in sources of dissolved organic matter. Our past data from this area showed that concentrations of TOC and TN increased from Everglades headwaters through the mangrove zone and then decrease with distance offshore. The high concentrations of TOC and TN in Florida Bay were due to a combination of terrestrial loading (Boyer and Jones, 1999), in situ production by seagrass and phytoplankton, and evaporative concentration (Fourqurean et al. 1993, Boyer et al. 1997).

Advection of Shelf and Florida Bay waters through the Sluiceway and passes accounted for this region and the inshore area of the Middle Keys as having highest TOC and TN of the FKNMS. In fact, isolated high concentrations in Fig 22 and 23 correspond to the location of SHORE stations. Strong offshore gradients in TOC and TN existed for all mainland Keys segments. The higher concentrations of TOC and TON in the inshore waters of the Keys imply a terrestrial source (anthropogenic) rather than simply benthic production and sediment re-suspension. Main Keys reef tract concentrations of TOC and TON were consistently the lowest in the FKNMS.




Figure 22. Distributions of surface total organic carbon across the region during 2015.




Figure 23. Distributions of surface total nitrogen across the region during 2015.

Much emphasis has been placed on assessing the impact of episodic phytoplankton blooms in Florida Bay on the offshore reef tract environment. In the past, spatial patterns of CHLA concentrations showed that the Shelf, NW Florida Bay, and the Ten Thousand Islands exhibited high levels of CHLA relative to the FKNMS. Also it showed that CHLA concentrations were typically higher in the Marquesas than in other areas of the FKNMS. When examined in context with the whole South Florida ecosystem, it is obvious that the Marquesas zone should be considered a continuum of the Shelf rather than a separate management entity. This shallow sandy area (often called the Quicksands) acts as a physical mixing zone between the Shelf and the Atlantic Ocean and is a highly productive area for other biota as well as it encompasses the historically rich Tortugas shrimping grounds. A CHLA concentration of $2 \mu\text{g l}^{-1}$ in the water column of a reef tract might be considered an indication of eutrophication. Conversely, a similar CHLA level in the Quicksands indicates a productive ecosystem which feeds a valuable shrimp fishery. In 2015 highest CHLA values preferentially occurred in the BKS and BKB and Marquesas areas, suggesting an important contribution from the Shelf (Fig. 24).

The oceanside transects in the Upper Keys exhibited the lowest overall CHLA concentrations of any area in the FKNMS. Transects off the Middle and Lower Keys showed that a drop in CHLA occurred at reef tract sites; there was no linear decline with distance from shore. Interestingly, CHLA concentrations in the Tortugas transect showed a similar pattern as the mainland Keys (Boyer and Briceño 2007). Inshore and Hawk Channel CHLA concentrations among Middle Keys, Lower Keys and Tortugas sites were not significantly different. As inshore CHLA concentrations in the Tortugas were similar to those in the Middle and Lower Keys, we see no evidence of persistent phytoplankton bloom transport from Florida Bay. The recently installed SHORE stations show higher CHLA concentrations than those of LK, MD, UK and OFF stations underscoring the anthropogenic impact.




Figure 24. Distributions of surface chlorophyll a across the region during 2015.

Along with TP, turbidity is probably the second most important determinant of local ecosystem health (Fig. 25). The fine grained, low density carbonate sediments in this area are easily re-suspended, rapidly transported, and have high light scattering potential. Sustained high turbidity of the water column indirectly affects benthic community structure by decreasing light penetration, promoting seagrass extinction. Large scale observations of turbidity clearly show patterns of onshore-offshore gradients which extend out onto the Shelf to the Marquesas (Stumpf et al. 1999). Strong turbidity gradients have been observed on the Shelf but reef tract levels remain remarkably low regardless of inshore levels. Elevated turbidity in the backcountry is most probably due to the shallow water column being easily re-suspended by wind and wave action.

Light extinction (K_d) was highest alongshore and improved with distance from land. This trend was expected as light extinction is related to water turbidity (Fig 26). However, in Keys waters, CDOM may be a more prominent driver of light penetration. For 2015, highest K_d was observed consistently in the Backcountry and nearshore areas.




Figure 25. Distributions of surface turbidity across the region during 2015.




Figure 26. Distributions of Light extinction across the region during 2015.

Surface SiO₂ concentrations exhibited a pattern similar to salinity (Fig. 27). The source of SiO₂ in this geologic area of carbonate rock and sediments is from siliceous periphyton (diatoms) growing in the Shark River Slough, Taylor Slough, and C-111 basin watersheds. Unlike the Mississippi River plume with CHLA concentrations of 76 μg l⁻¹ (Nelson and Dortch 1996), phytoplankton biomass on the Shelf (1-2 μg l⁻¹ CHLA) was not sufficient to account for the depletion of SiO₂ in this area. Therefore, SiO₂ concentrations on the Shelf are depleted mostly by mixing (although we no longer have data from the Shelf), allowing SiO₂ to be used as a semi-conservative tracer of freshwater in this system (Ryther et al. 1967; Moore et al. 1986).

In the Lower and Middle Keys, it is clear that the source of SiO₂ to the nearshore Atlantic waters is through the Sluiceway and Backcountry (Fig. 27). SiO₂ concentrations near the coast were elevated relative to the reef tract with much higher concentrations occurring in the Lower and Middle Keys than the Upper Keys. Our previous work indicates a significant relationship between SiO₂ concentrations in the Sluiceways and Florida Bay. There is an interesting peak in SiO₂ concentration in an area of the Sluiceway, which is densely covered with the seagrass, *Syringodium* (Fourqurean et al. 2002). We are unsure as to the source but postulate that it may be due to benthic flux.

The TN:TP ratio has been used as a relatively simple method of estimating potential nutrient limitation status of phytoplankton (Redfield 1967). Most of the South Florida hydroscape has TN:TP values >> 16:1, indicating the potential for phytoplankton to be limited by P at these sites (Fig. 28). However, most of the TN is not available to phytoplankton while much of the TP is labile. Therefore, using the TN:TP ratio overestimates potential P limitation and should be recognized as such.

Most of the FKNMS is routinely P limited using this metric. Phosphorous limitation in 2015 varies significantly with the seasons (Fig 28). During winter the highest limitation occurs in BKS, BKB and portions of UK; during spring limitation is milder overall; summer time strong limitation occurs in Marquesas and LK and extending into BKB and BKS; finally, during the fall limitation prevails in the Sluiceways and UK.




Figure 27. Distributions of surface silicate across the region during 2015.




Figure 28. Distributions of surface TN:TP ratio across the region during FY2015.

6. Numeric Nutrient Criteria Development

In order to gain a better understanding of the spatial patterns of water quality of the FKNMS, we attempted to reduce the complicated data matrix into fewer elements which would provide robust estimates of condition and connection. To this end we developed an objective classification analysis procedure which grouped stations according to water quality similarity (Briceño et al. 2013, Fig. 29).




Figure 29. Map of FKNMS showing segments derived from Factor and Cluster Analysis of biogeochemical data: OFF=Offshore; MAR=Marquesas; BKS=Back Shelf; BKB= Back Bay; LK= Lower Keys; MK= Middle Keys; UK= Upper Keys

This segmentation was proposed to EPA and FDEP for use in developing numeric nutrient criteria for FL estuaries and coastal waters. The approach was adopted by FDEP as Eight Maps of FL Marine Nutrient Regions, <https://www.flrules.org/Gateway/reference.asp?No=Ref-01215>, dated Oct. 19, 2011. The nutrient criteria for this region was subsequently developed using data from this Florida Keys Water Quality Monitoring Project and submitted to EPA in March 2013 as Implementation of FL Numeric Nutrient Standards, [NNC Implementation 3-11-13.pdf](#). On June 27, 2013, EPA approved [062713 epa approval nnc implementation document.pdf](#), FDEP Rule 62.300 (19) F.A.C. which includes 62.302.532 Estuary-Specific Numeric Interpretations of the Narrative Nutrient Criterion (the Coastal Estuary Rule), [62 302 final.pdf](#).

We believe that this accomplishment is an important achievement for a Federally-funded, University-operated water quality monitoring program and should be a model for future projects.

The large scale of this monitoring program allowed us to assemble a much more holistic view of broad physical/chemical/biological interactions occurring over the South Florida hydroscape. This confirms that rather than thinking of water quality monitoring as being a static, non-scientific pursuit it should be viewed as a tool for answering management questions and developing new scientific hypotheses.

We continue to maintain a website (<http://serc.fiu.edu/wqmnetwork/>) where data and reports from the FKNMS is integrated with the other parts of the SERC water quality network (Florida Bay, Whitewater Bay, Biscayne Bay, Ten Thousand Islands, and SW Florida Shelf) are available.

6.1. Acknowledgments

We thank all the field and laboratory technicians involved with this project, especially Jeff Absten, James Duquesnel, Mark Kershaw, Pete Lorenzo, Alexandra Serna Salazar, Sandro Stumpf, Milagros Timiraos, Antuanet Cuello, Diana Johnson, Ruth Justiniano, Grisel Menendez, and Dania Sancho. This project was possible due to continued funding by the FIU/US-EPA Agreement #X7 00D02412. This is Technical Report #799T of the Southeast Environmental Research Center at Florida International University.

7. References

- APHA. 1995. Automated method for molybdate-reactive silica. In A. D. Eaton, L. S. Clesceri, and A. E. Greenberg (Eds.), *Standard Methods for the Examination of Water and Wastewater*
- ALLEMAN, R. W., ET AL. 1995. Biscayne Bay surface water improvement and management. Technical supporting document. South Florida Water Management District.
- BOYER, J. N. 2006. Shifting N and P limitation along a north-south gradient of mangrove estuaries in South Florida. *Hydrobiologia* 269: 167-177.
- BOYER, J. N. AND H. O. BRICEÑO. 2007. FY2006 Annual Report of the South Florida Coastal Water Quality Monitoring Network. SFWMD/SERC Cooperative Agreement #4600000352. SERC Tech. Rep. T-351. [2006 CWQMN.pdf](#).
- BOYER, J. N. AND H. O. BRICEÑO. 2011. 2010 Annual Report of the Water Quality Monitoring Project for the Water Quality Protection Program of the Florida Keys National Marine Sanctuary. US EPA/FIU Agreement #X7-96410604-6. SERC Tech. Rep. T-536
- BOYER, J. N., J. W. FOURQUIREAN, AND R. D. JONES. 1997. Spatial characterization of water quality in Florida Bay and Whitewater Bay by multivariate analysis: Zones of similar influence (ZSI). *Estuaries* 20: 743-758.
- BOYER, J. N., J. W. FOURQUIREAN, AND R. D. JONES. 1999. Seasonal and long-term trends in water quality of Florida Bay (1989-1997). *Estuaries* 22:417-430.
- BOYER, J. N., AND R. D. JONES. 1999. Effects of freshwater inputs and loading of phosphorus and nitrogen on the water quality of Eastern Florida Bay, p. 545-561. In K. R. Reddy, G. A. O'Connor, and C. L. Schelske (eds.) *Phosphorus biogeochemistry in sub-tropical ecosystems*. CRC/Lewis Publishers, Boca Raton, Florida.
- BOYER, J. N., AND R. D. JONES. 2002. A view from the bridge: External and internal forces affecting the ambient water quality of the Florida Keys National Marine Sanctuary, p. 609-628. In J. W. Porter and K. G. Porter (eds.), *The Everglades, Florida Bay, and Coral Reefs of the Florida Keys: An Ecosystem Sourcebook*. CRC Press.
- BOYER, J. N., P. STERLING, AND R. D. JONES. 2000. Maximizing information from a water quality monitoring network through visualization techniques. *Estuarine, Coastal and Shelf Science* 50: 39-48.
- BRICEÑO, H. O., AND J. N. BOYER. 2010. Climatic controls on nutrients and phytoplankton biomass in a sub-tropical estuary, Florida Bay, USA. *Estuaries and Coasts* 33: 541-553.
- BRICEÑO, HENRY, JOSEPH BOYER AND PETER HARLEM. 2010. Proposed Methodology for the Assessment of Protective Numeric Nutrient Criteria for South Florida Estuaries and Coastal Waters. White paper submitted to Environmental Protection Agency Science Advisory Board. Dec 6 2010. FIU/SERC Contribution # T-501
- BRICEÑO, H.O. J.N. BOYER, J. CASTRO, AND P. HARLEM. 2013. Biogeochemical classification of south Florida's estuarine and coastal waters. *Marine Pollution Bulletin* 75: 187-204.
- BRICEÑO, H., R. GARCIA, P. GARDINALI, K. BOSWELL, A. SERNA AND E. SHINN. 2015. Design and implementation of dye-tracer injection test, Cudjoe Key, Florida Keys. FINAL REPORT.

- Submitted to CH2M Hills on behalf of Florida Keys Aqueduct Authority. FIU/SERC TR# T-723. 68 p
- CACCIA, V. G., AND J. N. BOYER. 2005. Spatial patterning of water quality in Biscayne Bay, Florida as a function of land use and water management. Marine Pollution Bulletin 50: 1416-1429.
- CACCIA, V. G., AND J. N. BOYER. 2007. A nutrient loading budget for Biscayne Bay, Florida. Marine Pollution Bulletin 54: 994–1008.
- CAPONE, D. G., AND B. F. TAYLOR. 1980. Microbial nitrogen cycling in a seagrass community, p. 153-161. *In* V. S. Kennedy (ed.), *Estuarine Perspectives*. Academic.
- CHRISTIAN, R. R., J. N. BOYER, D. W. STANLEY, AND W. M. RIZZO. 1991. Multi-year distribution patterns of nutrients in the Neuse River Estuary, North Carolina. Marine Ecology Progress Series 71:259-274.
- CORBETT, D. R., K. DILLON, W. BURNETT, AND J. CHANTON. 2000. Estimating the groundwater contribution into Florida Bay via natural tracers ^{222}Rn and CH_4 . Limnology and Oceanography 45:1546-1557.
- EPA. 1979. Handbook for Analytical Quality Control in Water and Wastewater Laboratories. EPA 600/4-79-019. Environmental Protection Agency, Environmental Monitoring and Support Laboratory, Cincinnati, OH.
- EPA. 1993. Water Quality Protection Program for the Florida Keys national Marine Sanctuary: Phase II Report. Battelle Ocean Sciences, Duxbury, MA and Continental Shelf Associates, Inc., Jupiter, FL.
- EPA. 1995. Water quality protection program for the Florida Keys National Marine Sanctuary: Phase III report. Final report submitted to the Environmental Protection Agency under Work Assignment 1, Contract No. 68-C2-0134. Battelle Ocean Sciences, Duxbury, MA and Continental Shelf Associates, Inc., Jupiter FL.
- EPA- Region 2. 1997. Non-Detect Policy. CENAN-OP-SD 28 February 1997
- FRANKOVICH, T. A., AND R. D. JONES. 1998. A rapid, precise, and sensitive method for the determination of total nitrogen in natural waters. *Mar. Chem.* 60:227-234.
- FOURQUREAN, J.W., M.D. DURAKO, M.O. HALL AND L.N. HEFTY. 2002. Seagrass distribution in south Florida: a multi-agency coordinated monitoring program, p. 497-522. *In* J. W. Porter and K. G. Porter (eds.), *The Everglades, Florida Bay, and Coral Reefs of the Florida Keys: An Ecosystem Sourcebook*. CRC Press.
- FOURQUREAN, J. W., R. D. JONES, AND J. C. ZIEMAN. 1993. Processes influencing water column nutrient characteristics and phosphorus limitation of phytoplankton biomass in Florida Bay, FL, USA: Inferences from spatial distributions. Estuarine, Coastal and Shelf Science 36:295-314.
- GIBSON, P. J., J. N. BOYER, AND N. P. SMITH. 2008. Nutrient Mass Flux between Florida Bay and the Florida Keys National Marine Sanctuary. Estuaries and Coasts 31: 21–32.
- GRASSHOFF, K. 1983a. Determination of nitrate. *In* K. Grasshoff, M. Erhardt, and K. Kremeling (Eds.), *Methods of Seawater Analysis*. Verlag Chemie, Weinheim, Germany.

- GRASSHOFF, K. 1983b. Determination of nitrite. In K. Grasshoff, M. Erhardt, and K. Kremeling (Eds.), *Methods of Seawater Analysis*. Verlag Chemie, Weinheim, Germany.
- HELSEL, D. 2005. *Nondetects and Data Analysis: Statistics for Censored Environmental Data*. John Wiley and Sons, New York. 250 p
- HIRSCH, R. M., R. B. ALEXANDER, AND R. A. SMITH. 1991. Selection of methods for the detection and estimation of trends in water quality. *Water Resources Research* 27:803-813.
- ISAKS, E. H., AND R. M. SRIVASTAVA. 1989. *An Introduction to Applied Geostatistics*. Oxford Press, 561 pp.
- KAISER, H.F. 1958. The varimax criterion for analytic rotation in factor analysis". *Psychometrika* 23 (3)
- KLEIN, C. J., AND S. P. ORLANDO JR. 1994. A spatial framework for water-quality management in the Florida Keys National Marine Sanctuary. *Bulletin of Marine Science* 54: 1036-1044.
- Koroleff, F. 1983. Determination of ammonia. In K. Grasshoff, M. Erhardt, and K. Kremeling (Eds.), *Methods of Seawater Analysis*. Verlag Chemie, Weinheim, Germany.
- MURPHY, J., AND J. P. RILEY. 1962. A modified single solution method for the determination of phosphate in natural water. *Anal. Chim. Acta* 27: 31-36.
- LAPOINTE, B. E., AND M. W. CLARK. 1992. Nutrient inputs from the watershed and coastal eutrophication in the Florida Keys. *Estuaries* 15: 465-476.
- LAPOINTE, B. E., AND W. R. MATZIE. 1996. Effects of stormwater nutrient discharges on eutrophication processes in nearshore waters of the Florida Keys. *Estuaries* 19: 422-435.
- LEE, T. N., M. E. CLARKE, E. WILLIAMS, A. F. SZMANT, AND T. BERGER. 1994. Evolution of the Tortugas gyre and its influence on recruitment in the Florida Keys. *Bulletin of Marine Science* 54: 621-646.
- LEE, T. N., E. WILLIAMS, E. JOHNS, D. WILSON, AND N. P. SMITH. 2002. Transport processes linking South Florida ecosystems, p. 309-342. *In* J. W. Porter and K. G. Porter (eds.), *The Everglades, Florida Bay, and Coral Reefs of the Florida Keys: An Ecosystem Sourcebook*. CRC Press.
- LEICHTER, J. J., S. R. WING, S. L. MILLER, AND M. W. DENNY. 1996. Pulsed delivery of subthermocline water to Conch Reef (Florida Keys) by internal tidal bores. *Limnology and Oceanography* 41: 1490-1501.
- LEICHTER, J. J. , AND S. L. MILLER. 1999. Predicting high-frequency upwelling: Spatial and temporal patterns of temperature anomalies on a Florida coral reef. *Continental Shelf Research* 19: 911-928.
- LEICHTER, J. J., H. L. STEWART, AND S. L. MILLER. 2003. Episodic nutrient transport to Florida coral reefs. *Limnology and Oceanography* 48:1394-1407.
- MOORE, W. S., J. L. SARMIENTO, AND R. M. KEY. 1986. Tracing the Amazon component of surface Atlantic water using ²²⁸Ra, salinity, and silica. *Journal of Geophysical Research* 91: 2574-2580.

- NELSON, D. M., AND Q. DORTCH. 1996. Silicic acid depletion and silicon limitation in the plume of the Mississippi River: evidence from kinetic studies in spring and summer. Marine Ecology Progress Series 136: 163-178.
- NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION. 1995. Florida Keys National Marine Sanctuary Draft Management Plan/Environmental Impact Statement.
- OVERLAND, J. E. AND R. W. PREISENDORFER. 1982. A significance test for principal components applied to cyclone climatology. Monthly Weather Review 110:1-4.
- PAUL J. H., ROSE J. B., BROWN J., SHINN E. A., MILLER S. AND FARRAH S. R. 1995. Viral tracer studies indicate contamination of marine waters by sewage disposal practices in Key Largo, Florida. Appl. Environ. Microbiol.61,2230-2234.
- PAUL, J.H., J.B. ROSE, S.C. JIANG, X. ZHOU, P. COCHRAN, C. KELLOGG, J.B. KANG, D. GRIFFIN, S. FARRAH AND J. LUKASIK. 1997. Evidence for groundwater and surface marine water contamination by waste disposal wells in the Florida Keys. Wat. Res. Vol. 31, No. 6, pp.
- PITTS, P. A. 1997. An investigation of tidal and nontidal current patterns in Western Hawk Channel, Florida Keys. Continental Shelf Research 17: 1679-1687.
- REDFIELD, A. C. 1958. The biological control of chemical factors in the environment. American Scientist 46: 205-222.
- REICH, C., E.A. SHINN, C. HICKEY AND A.B. TIHANSKY. 2001. Tidal and Meteorological Influences on Shallow Marine Groundwater Flow in the Upper Florida Keys in J. Porter and K.C Porter (Editors) The Everglades, Florida Bay, and Coral reefs of the Florida Keys. An Ecosystem Handbook. CRC Press. 1022 p.
- RUDNICK, D., Z. CHEN, D. CHILDERS, T. FONTAINE, AND J. N. BOYER. 1999. Phosphorus and nitrogen inputs to Florida Bay: the importance of the Everglades watershed. Estuaries 22: 398-416.
- RYTHER, J. H., D. W. MENZE, AND N. CORWIN. 1967. Influence of the Amazon River outflow on the ecology of the western tropical Atlantic, I. Hydrography and nutrient chemistry. Journal of Marine Research 25: 69-83.
- Shinn, E.A., C. Reich, D. Hickey and A.B. Tihansky. 1999a. Determination of Groundwater-Flow Direction and Rate Beneath Florida Bay, the Florida Keys and Reef Tract. http://sofia.usgs.gov/projects/index.php?project_url=grndwtr_flow Downloaded Oct 2014
- Shinn, E.A., R.S. Reese and C.D. Reich. 1999b. Fate and Pathways of Injection-Well Effluent in the Florida Keys. <http://sofia.usgs.gov/publications/ofr/94-276/index.html> Downloaded Oct 2014
- SMITH, N. P. 1994. Long-term Gulf-to-Atlantic transport through tidal channels in the Florida Keys. Bulletin of Marine Science 54: 602-609.
- SOLÓRZANO, L., AND J. H. SHARP. 1980. Determination of total dissolved phosphorus and particulate phosphorus in natural waters. Limnology and Oceanography 25:754-758.
- STUMPF, R. P., M. L. FRAYER, M. J. DURAKO, AND J. C. BROCK. 1999. Variations in water clarity and bottom albedo in Florida Bay from 1985-1997. Estuaries 22: 431-444.
- SZMANT, A. M., AND A. FORRESTER. 1996. Water column and sediment nitrogen and phosphorus distribution patterns in the Florida Keys, USA. Coral Reefs 15: 21-41.

WALSH, T. W. 1989. Total dissolved nitrogen in seawater: a new high temperature combustion method and a comparison with photo-oxidation. *Mar. Chem.* 26: 295-311.

YENTSCH, C. S., AND D. W. MENZEL. 1963. A method for determination of phytoplankton chlorophyll and phaeophytin by fluorescence. *Deep Sea Res.* 10: 221-231.